

String Theory and Black Holes

A thesis submitted to the
University of Mumbai
for the
Ph.D. Degree in Physics

by
Justin Raj David
Department of Theoretical Physics
Tata Institute of Fundamental Research
Mumbai 400 005, India.

1999

Abstract

This thesis aims to make precise the microscopic understanding of Hawking radiation from the D1/D5 black hole. We present an explicit construction of all the shortmultiplets of the $\mathcal{N} = (4, 4)$ SCFT on the symmetric product $\tilde{T}^4/S(Q_1Q_5)$. An investigation of the symmetries of this SCFT enables us to make a one-to-one correspondence between the supergravity moduli and the marginal operators of the SCFT. We analyse the gauge theory dynamics of the splitting of the D1/D5 system into subsystems and show that it agrees with supergravity. We have shown that the fixed scalars of the D1/D5 system couple only to $(2, 2)$ operators thus removing earlier discrepancies between D-brane calculations and semiclassical calculations. The absorption cross-section of the minimal scalars is determined from first principles upto a proportionality constant. We show that the absorption cross-section of the minimal scalars computed in supergravity and the SCFT is independent of the moduli.

Acknowledgements

Firstly, I would like to thank Spenta Wadia, my thesis advisor for inspiration, support and encouragement throughout my five years at TIFR. It is from him I learnt how to approach problems in physics. He has emphasized to me the importance of thinking from first principles and attacking physically relevant problems. During my early years in TIFR, Spenta's guidance helped me overcome the maze of superfluous literature and to concentrate on the basics. He emphasized to me the importance of independent thinking in physics. I thank him for sharing with me his wonderful ideas. His inspiring confidence helped me overcome moments of despair when all means of attacking a physics problem seemed to fail.

During the course of my five years I worked very closely with Gautam Mandal and Avinash Dhar. I have learnt (not completely) from Gautam the art of reducing a physics problem to its very essentials. It is amazing to see how simple a seemingly complicated problem is, when reformulated by Gautam. Avinash helped me to realize the importance of being persistent and pursuing a physics problem to its entire logical conclusion. Avinash and I were closely working on a problem which would not have yielded results had it not been for the constant encouragement and persistence of Avinash.

I owe most of my understanding of gauge theories to the wonderful course that Sumit Das took great pains to teach. The course stretched for the entire year of 1995. I thank the string community in TIFR for being active in fostering the nascent enthusiasm for the subject.

There are other members of the Theory group working in different fields who provided inspiration, help and guidance indirectly and unknowingly. They did it through their lectures, seminars and discussions. I have benefitted in this way from Deepak Dhar and Mustansir Barma.

The students of the Theory group during the past five years contributed a lot to help me imbibe the spirit of search and discovery. I would like to mention Abhishek Dhar whose uncompromising attitude towards physics continues to be a source of inspiration.

I would like to acknowledge the help from the efficient and friendly staff of the Department of Theoretical Physics.

Finally I would like to thank my family and friends for their support in helping me pursue physics as a career.

Contents

1	Introduction	1
1.1	Black hole physics	1
1.2	Black holes in string theory	4
1.3	The D1/D5 black hole	6
1.4	Outline of the Thesis	11
2	The Microscopic Theory of the D1/D5 system	20
2.1	The gauge theory of the D1/D5 system	20
2.2	The instanton moduli space	25
2.3	The SCFT on the orbifold $\mathcal{M}$	27
2.3.1	The $\mathcal{N} = 4$ superconformal algebra	27
2.3.2	Free field realization of $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$	28
2.3.3	The $SO(4)$ algebra	29
2.3.4	The supergroup $SU(1, 1 2)$	31
2.4	Short multiplets of $SU(1, 1 2)$	31
2.5	The resolutions of the symmetric product	32
2.5.1	The untwisted sector	32
2.5.2	Z_2 twists.	33
2.5.3	Higher twists	36
2.6	The chiral primaries of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$	37
2.6.1	The k-cycle twist operator	37
2.6.2	The complete set of chiral primaries	39
2.7	Shortmultiplets of $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$	42

2.8	The location of the symmetric product	43
2.9	The linear sigma model	45
2.9.1	The linear sigma model description of R^4/Z_2	45
2.9.2	The gauge theory description of the moduli of the D1/D5 system . .	48
2.9.3	The case $(Q_1, Q_5) \rightarrow (Q_1 - 1, Q_5) + (1, 0)$: splitting of 1 D1-brane . .	52
2.9.4	The dynamics of the splitting at the singularity of the Higgs branch .	53
3	Coupling with the bulk fields	56
3.1	Near horizon symmetry	56
3.2	Classification of the supergravity modes	58
3.2.1	Comparison of supergravity shortmultiplets with SCFT	61
3.3	The supergravity moduli	61
3.4	AdS ₃ /CFT ₂ correspondence	63
3.5	Supergravity moduli and the marginal operators	63
3.6	Fixed scalars	65
3.7	Intermediate Scalars	67
3.8	Supergravity from gauge theory	67
4	The Hawking Process	73
4.1	Supergravity calculation of absorption/Hawking radiation in presence of moduli	73
4.2	Near horizon geometry of the D1/D5 black hole	74
4.3	The coupling with the bulk fields in the Ramond sector	77
4.4	Determination of the strength of the coupling μ	77
4.4.1	Evaluation of the tree-level vertices in supergravity	80
4.4.2	Two-point function	82
4.5	The black hole state	83
4.6	Absorption cross-section as thermal Green's function	86
4.7	Absorption cross-section of minimal scalars from the D1/D5 SCFT	88
4.8	Independence of Hawking radiation on D1/D5 moduli	89
4.9	Entropy and area	94

Chapter 1

Introduction

A unified theory of fundamental interactions should include within its framework a quantum theory of gravity. The quantization of Einstein's theory of gravity using conventional methods poses a host of problems. Einstein's theory of gravity is a non-renormalizable field theory. Therefore one can not extract meaningful answers from quantum perturbation theory. The existence of black holes as solutions in Einstein's theory of gravity is another stumbling block. Arguments involving physics of black holes within the framework of quantum mechanics seem to lead to the conclusion that quantum mechanical evolution of black hole is not unitary. This contradicts the basic rules of quantum mechanics.

At present string theory is the leading candidate for a unified theory of fundamental interactions. The massless spectrum of string theory includes the graviton. The low energy effective action of string theory includes the Einstein action. String theory is also consistent with the rules of quantum mechanics. Thus the natural generalization of Einstein's theory within the framework of quantum mechanics is string theory. Furthermore string theory is perturbatively finite. This cures the non-renormalizability of gravity. The understanding of non-perturbative spectra of string theory like D-branes has paved the way for addressing the problems of black hole physics. In this thesis we will attempt to make precise the description of black hole physics within the framework of string theory.

1.1 Black hole physics

Let us briefly review some general properties of black holes. [1] Black holes are objects which result as end points of gravitational collapse of matter. For masses greater than 3.6 solar mass, the gravitational force overcomes all other forces and the matter generically collapses

into a black hole (in some exceptional cases a naked singularity might result). This would suggest that to specify a black hole it is necessary to give in detail the initial conditions of the collapse. As we will see below a black hole is completely specified by a few parameters only.

To introduce various concepts related to black hole we will discuss two examples of black holes. First, let us consider the Schwarzschild black hole in $3 + 1$ dimensions. It is a time independent, spherically symmetric solution of pure Einstein gravity. Its metric is given by

$$ds^2 = - \left(1 - \frac{2G_N M}{r}\right) dt^2 + \left(1 - \frac{2G_N M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2 \quad (1.1)$$

where t refers to time, r the radial distance, Ω the solid angle in 3 dimensions and G_N the Newton's constant. We have chosen units so that the velocity of light, $c = 1$. The surface $r = 2G_N M$ is called the event horizon. It is a co-ordinate singularity but not a curvature singularity. Light-like geodesics and time-like geodesics starting at $r < 2G_N M$ end up at $r = 0$ (the curvature singularity) in finite proper time. This means that classically the black hole is truly black, it cannot emit anything. Note that the solution is completely specified by only one parameter M , the mass of the black hole.

Next we consider the Reissner-Nördstrom black hole. It is a time independent, spherically symmetric solution of Einstein gravity coupled to the electromagnetic field. The solution is given by the following backgrounds.

$$\begin{aligned} ds^2 &= - \left(1 - \frac{2G_N M}{r} + \frac{G_N Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2G_N M}{r} + \frac{G_N Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2, \\ A_0 &= \frac{Q}{r}. \end{aligned} \quad (1.2)$$

where A_0 is the time component of the vector potential. This solution carries a charge Q . There are two co-ordinate singularities at

$$r_+ = G_N M + \sqrt{G_N^2 M^2 - G_N Q^2} \quad (1.3)$$

and

$$r_- = G_N M - \sqrt{G_N^2 M^2 - G_N Q^2} \quad (1.4)$$

The event horizon is at $r = r_+$. When $M = |Q|/\sqrt{G_N}$, the outer horizon at $r = r_+$ and the inner horizon at $r = r_-$ coincide. A black hole with coincident inner and outer horizons is

called an extremal black hole. Note that in this case the black hole is completely specified by its mass M and the charge Q .

In general, collapsing matter results in black holes which are completely specified by the mass M , the $U(1)$ charges Q_i and the angular momentum J . This is called the no hair theorem. Whatever other information (for example, multipole moments) present decays exponentially fast during the collapse. Thus, all detailed information carried by the collapsing matter is completely lost.

So far we have discussed the black holes only classically. In the seventies, the works of Bekenstein, Hawking and others furthered the understanding of black holes within the framework of quantum mechanics. It was found by Hawking that the Schwarzschild black hole is not truly black. A semi-classical calculation by Hawking showed that it emits radiation with the spectrum of a black body at a temperature T given by

$$T = \frac{\hbar}{8\pi G_N M} \quad (1.5)$$

The quantum nature of this effect is clearly evident from the fact that the temperature is proportional to $\hbar$. For the Reissner-Nördstrom black hole the temperature of Hawking radiation is

$$T = \frac{(r_+ - r_-)\hbar}{4\pi r_+^2} \quad (1.6)$$

One notes that the extremal Reissner-Nördstrom black hole does not Hawking radiate. In general the Hawking temperature turns out to be function of mass, the charges and the angular momentum alone. Thus even semi-classical effects do not provide further information of the black hole. The works in the seventies culminated in the following laws of black holes which are analogous to the laws of thermodynamics.

i. First Law: Two neighboring black hole equilibrium states are related by

$$dM = Td\left[\frac{A}{4G_N\hbar}\right] + \Phi_i dQ_i + \Omega dJ \quad (1.7)$$

where A is the area of the event horizon, Φ the electric surface potential and Ω the angular velocity. For the special case of the Reissner-Nördstrom black hole the first law reduces to

$$dM = Td\left[\frac{A}{4G_N\hbar}\right] + \Phi dQ. \quad (1.8)$$

where $\Phi = Q/r_+$.

ii. Second Law: Black holes have entropy S given by

$$S = \frac{A}{4G_N\hbar} \tag{1.9}$$

In any irreversible process, the sum of the entropy of the black hole and the surroundings always increases.

The above facts taken together lead to the following puzzle. Consider a black hole formed out of the collapse of a pure state. It eventually evaporates through Hawking radiation. The radiation is purely thermal. The final state is a mixed state. Thus we have a situation in which a pure state has evolved into a mixed state. This contradicts the unitary evolution of quantum mechanics.

To emphasize where we lack in understanding let us contrast the above situation with that of radiation from a black body. A black body made up of a pure state radiates thermally. This is because we have averaged over the various micro-states of the black body. The radiation would contain sufficient information to reconstruct the pure state of the black body. In this case we do not have a contradiction with quantum mechanics.

The difficulty with the understanding of black holes within the framework of quantum mechanics is because we do not know what are the microstates which contribute to the entropy of the black hole. Any serious candidate for the quantum theory of gravity should enable us to understand black holes from a microscopic point of view. It should provide us with a microscopic theory analogous to the way statistical mechanics is the microscopic theory for thermodynamics. String theory has demonstrated the potential to fulfill this requirement.

1.2 Black holes in string theory

The low energy effective action of string theory has black holes as solutions of its equation of motion. Based on earlier suggestions [2, 3, 4, 5] an attempt was made to understand the entropy of black holes in string theory. [6, 7] showed that the microscopic states contributing to the entropy of certain black holes in string theory arises due to the degeneracy of the

perturbative string spectrum. For bosonic string theory it was seen [6] that a very massive string with mass M at strong string coupling g_s has a Schwarzschild radius, $r_{\text{Schwarzschild}} \sim g_s^2 M$ larger than the string length $\sqrt{\alpha'}$ and therefore can be treated as a black hole. The entropy of states of the perturbative string for a given mass level can be computed as follows. One evaluates the degeneracy of the string states at a given mass. The logarithm of this degeneracy is the entropy of the string states at that mass level. For a perturbative string of mass M , the entropy is proportional to M . But it is well known that the entropy of a Schwarzschild black hole is proportional to M^2 . It was hoped that strong coupling effects would renormalize the mass of the string so that the entropy of the string could agree with that of the corresponding black hole. If one has a situation where mass does not get renormalized then the entropy evaluated at weak coupling should agree with that calculated at strong coupling. In superstring theory there are states whose masses are not renormalized. These are the BPS states. It was shown in [7] that a massive BPS string of heterotic string theory compactified on T^6 has an entropy due to perturbative string states which is proportional to the area of the stretched horizon. The supergravity configuration corresponding to this BPS string has a horizon of zero area. Thus it was necessary to find a black hole solution in string theory which is BPS and has finite area of horizon. Such black holes are known, but they carry Ramond-Ramond charges. They are solitonic solutions of the supergravity equations of motion. It was not until the discovery of the carriers of these charges known as D-branes [8] and their understanding in terms of perturbative string theory that a microscopic understanding of black holes developed. Strominger and Vafa in [9] calculated the entropy of an extremal and BPS black hole of type IIB string theory compactified on $K3 \times S^1$. This extremal black hole was obtained by wrapping D5-branes along the 4-cycle of $K3$ and introducing momentum along the S^1 . Strominger and Vafa showed that the entropy calculated as the logarithm of the degeneracy of states in the microscopic theory corresponding to the extremal black hole agrees precisely with that computed from supergravity.

1.3 The D1/D5 black hole

A black hole solution which is BPS and has a finite area of horizon was also studied by Callan and Maldacena [10]. It is a solution of type IIB string theory compactified on $T^4 \times S^1$ to five dimensions. This solution avoids the complication of having to deal with $K3$. This black hole will be the working model in this thesis ¹. We will discuss the solution in detail below.

The relevant terms in the supergravity action for the solution are

$$\frac{-1}{128\pi^7\alpha'^4 g_s^2} \int d^{10}x \sqrt{-g} \left[e^{-2\phi} (R - 4(d\phi)^2 + \frac{1}{12}(dB)^2) \right] \quad (1.10)$$

B denotes the Ramond-Ramond two form potential, ϕ the dilaton and g the metric. To construct the supergravity solution, we take Q_5 D5-branes and align them along the 5, 6, 7, 8, 9 directions and Q_1 D1-branes along the x_5 direction. We then compactify the 6, 7, 8, 9 directions on a torus T^4 of volume V_4 and the x_5 direction on a circle of radius R_5 . To obtain a black hole with nonzero area for its event horizon we must introduce N units of momentum along the x_5 direction. The supergravity solution in 10 dimensions for this configuration is obtained by setting all the other fields of type IIB supergravity to zero and setting the fields in the action to the following values.

$$\begin{aligned} ds^2 &= f_1^{-\frac{1}{2}} f_5^{-\frac{1}{2}} (-dt^2 + dx_5^2 + k(dt - dx_5)^2) + f_1^{\frac{1}{2}} f_5^{\frac{1}{2}} (dx_1^2 + \dots + dx_4^2) \\ &\quad + f_1^{\frac{1}{2}} f_5^{-\frac{1}{2}} (dx_6^2 + \dots + dx_9^2), \\ e^{-2\phi} &= \frac{1}{g_s^2} f_5 f_1^{-1}, \\ B_{05} &= \frac{1}{2} (f_1^{-1} - 1), \\ H_{abc} &= (dB)_{abc} = \frac{1}{2} \epsilon_{abcd} \partial_d f_5, \quad a, b, c, d = 1, 2, 3, 4 \end{aligned} \quad (1.11)$$

where f_1 , f_5 and k are given by

$$f_1 = 1 + \frac{16\pi^4 g_s \alpha'^3 Q_1}{V_4 r^2}, \quad f_5 = 1 + \frac{g_s \alpha' Q_5}{r^2}, \quad k = \frac{16\pi^4 g_s^2 \alpha'^3 N}{V_4 R_5^2 r^2} \quad (1.12)$$

Here $r^2 = x_1^2 + x_2^2 + x_3^2 + x_4^2$ denotes the distance measured in the transverse direction to all the D-branes. On compactifying the above solution to 5 dimensions using the Kaluza-Klein

¹In the rest of the thesis, we set $\hbar = 1$.

ansatz one obtains an extremal black hole with the horizon at $r = 0$ but with a finite area.

The entropy of the 5 dimensional black hole is given by

$$S = 2\pi\sqrt{Q_1 Q_5 N} \quad (1.13)$$

This solution preserves 4 supersymmetries out of the 32 supersymmetries of type IIB theory.

It is a BPS configuration.

A non-extremal version of the above black hole solution can be obtained by having Kaluza-Klein momentum N which is distributed along the left and right directions of x_5 . This breaks all the 32 supersymmetries. Such a solution in ten dimensions is given by the following background.

$$\begin{aligned} e^{-2\phi} &= \frac{1}{g_s^2} \left(1 + \frac{r_5^2}{r^2}\right) \left(1 + \frac{r_1^2}{r^2}\right)^{-1}, \\ H &= \frac{2r_5^2}{g_s} \epsilon_3 + 2g_s e^{-2\phi} r_1^2 *_6 \epsilon_3, \\ ds^2 &= \left(1 + \frac{r_1^2}{r^2}\right)^{-1/2} \left(1 + \frac{r_5^2}{r^2}\right)^{-1/2} \left[-dt^2 + dx_9^2\right. \\ &\quad \left.+ \frac{r_0^2}{r^2} (\cosh \sigma dt + \sinh \sigma dx_9)^2 + \left(1 + \frac{r_1^2}{r^2}\right) (dx_5^2 + \dots dx_8^2)\right] \\ &\quad + \left(1 + \frac{r_1^2}{r^2}\right)^{1/2} \left(1 + \frac{r_5^2}{r^2}\right)^{1/2} \left[\left(1 - \frac{r_0^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega_3^2\right], \end{aligned} \quad (1.14)$$

where $*_6$ is the Hodge dual in the six dimensions $x_0, \dots, x_5$ and ϵ_3 is the volume form on the unit three-sphere. x_5 is periodically identified with period $2\pi R_5$ and directions $x_6, \dots, x_9$ are compactified on a torus T^4 of volume V_4 . Ω_3 is the volume of the unit three-sphere in the transverse directions. This solution is parameterized by six independent quantities – $r_1, r_5, r_0, \sigma, R_5$ and V_4 . These are related to the number of D1-branes, D5-branes and Kaluza-Klein momentum on x_5 as follows.

$$\begin{aligned} Q_1 &= \frac{V_4}{64\pi^6 g_s^2 \alpha'^3} \int e^{2\phi} *_6 H = \frac{V_4 r_1^2}{16\pi^4 \alpha'^3 g_s}, \\ Q_5 &= \frac{1}{4\pi^2 \alpha'} \int H = \frac{r_5^2}{g_s \alpha'}, \\ N &= \frac{R_5^2 V_4 r_0^2}{32\pi^4 \alpha'^4 g_s^2} \sinh 2\sigma. \end{aligned} \quad (1.15)$$

On compactifying this solution to five dimensions using the Kaluza-Klein ansatz one obtains a five-dimensional black hole with a horizon at $r = r_0$. The entropy and the mass of this black hole is given by

$$\begin{aligned} S &= \frac{A}{4G_5} = \frac{2\pi^2 r_1 r_5 r_0 \cosh 2\sigma}{4G_5}, \\ M &= \frac{\pi}{4G_5} (r_1^2 + r_5^2 + \frac{r_0^2 \cosh 2\sigma}{2}), \end{aligned} \quad (1.16)$$

where the five-dimensional Newton's constant is

$$G_5 = \frac{4\pi^5 \alpha'^4 g_s^2}{V_4 R_5} \quad (1.17)$$

We will now discuss the various limits in which physics of this black hole solution can be understood [12]. As the supergravity action has a $1/g_s^2$ prefactor the classical limit can be obtained by setting $g_s \rightarrow 0$. The fields are held fixed in the classical limit. From (1.14) and (1.15) we see that this is equivalent to

$$g_s \rightarrow 0, \quad (1.18)$$

with $g_s Q_1$, $g_s Q_5$, $g_s^2 N$ fixed.

The formulae in (1.15) indicate that this is also equivalent to

$$g_s \rightarrow 0, \quad (1.19)$$

with r_1 , r_5 , r_N fixed.

where $r_N = r_0 \sinh \sigma$. Hawking's semi-classical analysis of the radiation emitted by the black hole is valid in this limit.

The D1/D5 black hole is a solution of the equations of motion of type IIB supergravity. Type IIB supergravity is the low energy effective theory of type IIB string theory. Therefore the classical limit (1.18) is described by genus zero closed string theory. In string theory for every genus there is an α' expansion. Closed string perturbation is valid when the typical length scales involved are large compared to the string length $\sqrt{\alpha'}$. From (1.14) we find the typical length scales involved are r_1, r_5, r_N . Thus the α' expansion is valid when r_1, r_5, r_N are large compared to string length. From (1.15) this is equivalent to

$$g_s Q_1 \gg 1, \quad g_s Q_5 \gg 1, \quad g_s^2 N \gg 1 \quad (1.20)$$

The D1/D5 black hole can also be described by the constituent D-branes. D-brane perturbation theory involves open string loops. Open string loops have factors of $g_s Q_1$ or $g_s Q_5$, corresponding to the fact that the open string loops can end on any of the D-branes. Hence the classical limit (1.18) corresponds to a large N limit of the open string field theory. Perturbation theory of the large N limit of open string theory is good if

$$g_s Q_1 \ll 1, g_s Q_5 \ll 1, g_s^2 N \ll 1. \quad (1.21)$$

In the limit (1.21) the classical solution is still a black hole, though the typical size of this solution in the limit (1.21) is smaller than the string length. This is because in the classical limit (1.18) the entropy S in (1.16) diverges. It costs an infinite amount of entropy for the black hole to lose any finite fraction of its mass in outgoing radiation. The second law thus prohibits radiation from escaping and black holes are black in the classical limit (1.21), independent of their size. The microscopic understanding of the D1/D5 black hole involves understanding the D1/D5 black hole in terms of the constituent D-branes. Thus any calculation of a macroscopic property of the black hole like entropy or Hawking radiation in terms of the D-branes is valid only in the small black hole region. Surprisingly many of the calculations done in the limit (1.21) agree with those done in the limit (1.20). The reason is generally ascribed to some non-renormalization theorems.

The entropy of the extremal and the near extremal black hole discussed above was derived in [13] using a phenomenologically motivated microscopic model. It consists of a string wound along the x_5 direction. The string was assumed to carry the effective degrees of freedom of the D1/D5 black hole. The length of the string is $Q_1 Q_5 R_5$. The string is assumed to have oscillation only along the directions of the torus T^4 . It has four bosonic and four fermionic degrees of freedom. For an extremal black hole with N units of Kaluza-Klein momentum along x_5 the string is excited to a level $N Q_1 Q_5$. This is necessary because of the quantization of the Kaluza-Klein momentum in units of $1/R_5$. The value of $L_0 - \bar{L}_0$ of the string is identified with the Kaluza-Klein momentum of the black hole. Thus we have

$$L_0 - \bar{L}_0 = \frac{N Q_1 Q_5}{Q_1 Q_5 R_5} = \frac{N}{R_5} \quad (1.22)$$

Using Cardy's formula,

$$S_{\text{string}} = 2\pi\sqrt{NQ_1Q_5} \quad (1.23)$$

This agrees with (1.13) obtained from the supergravity solution. We will call this phenomenological model the 'long string' model.

Hawking radiation of scalars from the near-extremal black hole was understood microscopically from the long string model in [14, 15]. To derive Hawking radiation it was necessary to couple the microscopic model to various fields of supergravity. In [14, 15] this was done using couplings derived from a Dirac-Born-Infeld (DBI) action of the long string. This method gave results which agreed with supergravity for minimal scalars. These scalars are massless in the black hole geometry. A detailed calculation by [12] showed that using the long string as the microscopic theory for the D1/D5 black hole reproduces the Hawking rate taking into account of the grey body factors of the supergravity. The Hawking rate with grey body factor and the decay rate using the long string is given by

$$\Gamma_H = \Gamma_{\text{string}} = 2\pi^2 r_1^2 r_5^2 \frac{\pi\omega}{2} \frac{1}{e^{\frac{\omega}{2T_L}} - 1} \frac{1}{e^{\frac{\omega}{2T_R}} - 1} \frac{d^4k}{(2\pi)^4} \quad (1.24)$$

where

$$\begin{aligned} T_L &= \frac{1}{\pi} \frac{r_o e^\sigma}{2r_1 r_5} = \frac{1}{\pi R_5 Q_1 Q_5} \sqrt{N_L} \\ T_R &= \frac{1}{\pi} \frac{r_o e^{-\sigma}}{2r_1 r_5} = \frac{1}{\pi R_5 Q_1 Q_5} \sqrt{N_R} \end{aligned} \quad (1.25)$$

The right movers of the string are excited to a level N_R and the left movers of the string are excited to a level N_L .

In spite of these achievements the long string model had many shortcomings. As we will see the long string model could explain Hawking radiation of those minimal scalars which correspond to the metric fluctuation of the torus T^4 . In the supergravity background of the D1/D5 black hole there are four additional minimal scalars discussed in Chapter 2 and Chapter 3. The long string model has no way of explaining Hawking radiation of these additional minimal scalars. The long string model failed to reproduce the Hawking rate expected in supergravity for the fixed scalars [16]. These scalars are massive in the black hole geometry. This suggests that the long string model along with the DBI couplings could

not be a microscopic model for the D1/D5 black hole. In some ways the model has an ad-hoc nature. It does not explain why the string has to be of length $Q_1 Q_5 R_5$. The DBI action is valid only for a single set of branes. Its use for the case of multiple D1, D5-branes is at best a crude approximation.

The first attempt to obtain a microscopic model for the D1/D5 black hole based on the low energy effective theory of the D1/D5 system was made in [17]. It was found that the low energy effective degrees of freedom turn out to be that of a effective string. The fact that the Kaluza-Klein momenta has to be quantized in units of $1/(R_5 Q_1 Q_5)$ was explained by constructing the effective string from the infrared gauge invariant degrees of freedom of the D1/D5 gauge theory. However using the methods of [17] it was difficult to obtain the couplings of the effective theory with the supergravity.

1.4 Outline of the Thesis

In this thesis we attempt to make precise the microscopic derivation of Hawking radiation for the D1/D5 black hole. Abstracting from the discussion in the last paragraphs of the previous section, the requirements for understanding Hawking radiation are: 1. Discovering the microscopic degrees of freedom responsible for the black hole entropy. 2. Deriving how these degrees of freedom couple with the fields present in the bulk of the space-time. 3. Explaining the process of the interaction of the bulk fields with the microscopic degrees of freedom to produce Hawking radiation within a unitary framework.

The outline of this thesis is as follows. In Chapter 2, we discuss the microscopic degrees of freedom relevant for the D1/D5 black hole. The microscopic degrees of freedom are best described by first isolating the degrees of freedom for a system closely related to the D1/D5 black hole. This system is obtained by type IIB theory compactified on T^4 with Q_5 D5-branes wrapped on T^4 . The compact directions are x_6, x_7, x_8, x_9 . The single non-compact direction of the D5-branes is chosen to be along the x_5 direction. The Q_1 D1-branes are also aligned along the x_5 direction. The difference between the D1/D5 black hole and this system is that the x_5 direction is not compact in the latter. The supergravity solution is that of a black string in six dimensions. This solution lifted to ten dimensions is given by

setting $N = 0$ in (1.11). We will call this system the D1/D5 system.

We will study the D1/D5 system from the D-brane point of view. As we have seen before this is valid in the limit (1.21). size. For understanding Hawking radiation, it is sufficient to study the low energy effective theory of the D1/D5 system. The low energy theory of the D1/D5 system is a $1 + 1$ dimensional supersymmetric gauge theory with gauge group $U(Q_1) \times U(Q_5)$. It has eight supersymmetries. The matter content of this theory consists of hypermultiplets transforming in the adjoint representation of each of the gauge groups and hypermultiplets transforming as bifundamentals of $U(Q_1) \times \overline{U(Q_5)}$. We will review in detail the field content of this theory and its symmetries. The bound state of the D1 and D5-branes is described by the Higgs branch of this theory. The Higgs branch in the infrared will flow to a certain $\mathcal{N} = (4, 4)$ superconformal field theory (SCFT). The D1-branes bound to the D5-branes can be considered as instanton strings of the $U(Q_5)$ gauge theory. From this point of view the target space of the SCFT is the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on T^4 . This moduli space is known to be a resolution of the orbifold $(\tilde{T}^4)^{Q_1 Q_5} / S(Q_1 Q_5)$ which we shall denote by $\mathcal{M}$. $\tilde{T}^4$ can be distinct from the compactification torus T^4 . The evidence for this is mainly topological and is related to dualities which map the black string corresponding to D1/D5 system to a perturbative string of Type IIB theory with Q_1 units of momentum and Q_5 units of winding along the x_5 direction. We will discuss in detail a realization of this orbifold SCFT as a free field theory with identifications. The symmetries of this SCFT including a new $SO(4)$ algebra will also be discussed.

In this thesis we will use the Higgs branch of the D1/D5 system, realized as a $\mathcal{N} = (4, 4)$ SCFT on a resolution of the orbifold $\mathcal{M}$, to provide a microscopic understanding of Hawking radiation from the D1/D5 black hole. The point of view adopted in this thesis is to attempt at a first principle derivation of Hawking radiation from the Higgs branch of the D1/D5 system. The long string model used by [13] is at best a phenomenologically motivated theory.

The complete specification of the D1/D5 system involves the specification of the various supergravity moduli. The supergravity solution (1.11) is the solution with no moduli. We would like to understand what the SCFT is corresponding to the D1/D5 system at generic values of the moduli. To this end we construct all the marginal operators of the $\mathcal{N} = (4, 4)$

SCFT on $\mathcal{M}$ including operators involving twist field which correspond to blowing up modes. We classify the marginal operators according to the short multiplets of the global part of the $\mathcal{N} = (4, 4)$ superalgebra and a new $SO(4)$ algebra. It is known from supergravity that the D1/D5 bound state (Q_1, Q_5) with no moduli turned on is marginally stable with respect to decay to subsystems (Q'_1, Q'_5) and (Q''_1, Q''_5) . Analyzing the gauge theory of the D1/D5 system we find that the dynamics of this splitting is described by an effective $(4, 4)$ $U(1)$ theory coupled to $Q'_1 Q''_5 + Q''_1 Q'_5$ hypermultiplets. We show, by an analysis of the D-term equations and the potential, that the splitting is possible only when the Fayet-Iliopoulos terms and the theta term of the effective gauge theory are zero [18].

In Chapter 3, we take the next step towards the microscopic understanding of Hawking radiation. We find the precise coupling of the fields of the supergravity to the microscopic SCFT. This is given by a specific SCFT operator $\mathcal{O}(z, \bar{z})$, which couples to the supergravity field ϕ in the form of an interaction

$$S_{\text{int}} = \mu \int d^2 z \phi(z, \bar{z}) \mathcal{O}(z, \bar{z}) \quad (1.26)$$

where μ is the strength of the coupling. As the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ is an “effective” theory of the D1/D5 system, it is difficult to fix the coupling of this theory to the supergravity fields. Traditionally couplings in effective theories have been fixed using the method of symmetries. For example, the pion-nucleon coupling in the pion model is fixed by symmetries. The pion Π^i transforms as a **3** of the $SU(2)$ isospin symmetry. The nucleon N transforms as a **2** under this $SU(2)$. Therefore the pion-nucleon coupling consistent with this symmetry is

$$\mathcal{L}_{\text{int}} = g_{\text{eff}} \bar{N} \sigma^i \Pi^i \gamma_5 N \quad (1.27)$$

where g_{eff} is the strength of the pion-nucleon coupling. The operator $\bar{N} \sigma^i \gamma_5 N$ transforms as **3** under the isospin $SU(2)$. The γ_5 occurs because the pion is a pseudoscalar. The strength of the pion-nucleon coupling g_{eff} can be fixed only by appealing to the QCD, the microscopic theory relevant for the pion-nucleon system. In this thesis we fix the operator in the SCFT corresponding to the supergravity field using the method of symmetries. It is seen that in the near horizon limit the D1/D5 system exhibits enhanced symmetries. This is a special case of the AdS/CFT correspondence. The near horizon geometry of the D1/D5 system reduces

to that of $AdS_3 \times S^3 \times T^4$. The AdS/CFT correspondence [19] states that string theory on $AdS_3 \times S^3 \times T^4$ is dual to the 1 + 1 dimensional SCFT describing the Higgs branch of the D1/D5 system on T^4 . The radius of S^3 is $2\pi\alpha'(g_s^2 Q_1 Q_5 / V_4)^{1/4}$. The Kaluza-Klein modes on T^4 is of the order of the string length. Thus for large $Q_1 Q_5$, type IIB string theory on $AdS_3 \times S^3 \times T^4$ passes over to 6 dimensional (2, 2) supergravity on $AdS_3 \times S^3$. We will work in the supergravity limit. The evidence for this correspondence comes from symmetries. The isometries of AdS_3 correspond to the global part of the Virasoro group of the SCFT. The R-symmetry of the SCFT is identified with the isometry of the S^3 . The number of supersymmetries of the bulk get enhanced to 16 from 8. These correspond to the global supersymmetries of the $\mathcal{N} = (4, 4)$ superalgebra of the SCFT. Thus the global part of the superalgebra of the SCFT is identified with the $AdS_3 \times S^3$ supergroup, $SU(1, 1|2) \times SU(1, 1|2)$.

Therefore a viable strategy to fix the coupling is to classify both the bulk fields and the SCFT operators according to the symmetries. The question then would be if this procedure can fix the SCFT operator required for analysing Hawking radiation. We will review the classification of the entire set of Kaluza-Klein modes of the six-dimensional supergravity on S^3 as short multiplets of $SU(1, 1|2) \times SU(1, 1|2)$ [20]. We use symmetries, including a new global $SO(4)$ algebra, to identify the marginal operators constructed in Chapter 2 with their corresponding supergravity fields. This enables us to identify the operators corresponding to the minimal scalars. We show that the four of the minimal scalars which are not metric fluctuations of T^4 correspond to the blow up modes of the orbifold $(\tilde{T}^4)^{Q_1 Q_5} / S(Q_1 Q_5)$. The long string model has no operators corresponding to these minimal scalars because it is not based on an orbifold. Therefore it cannot be the effective theory for the D1/D5 system. We also find the representation of the operators of the fixed scalars under the short multiplet of the supergroup $SU(1, 1|2) \times SU(1, 1|2)$ [18, 21]. We show that the SCFT operators corresponding to the all the fixed scalars have conformal dimension $(h, \bar{h}) = (2, 2)$. The long string along with the DBI action predicts that the fixed scalars also couple to operators with dimension (1, 3) and (3, 1). Hawking radiation computed from the long string model using (1, 3) and (3, 1) operators gave results which did not agree with that computed from supergravity. Operators of dimension (2, 2) gave results which agreed with supergravity. Thus

this method of fixing the coupling removes the discrepancy found earlier for the fixed scalars using the long string model. In Chapter 4 we will see that the Hawking rate/absorption cross-section is determined upto a proportionality constant by the dimension of the SCFT operator corresponding to the supergravity field. Thus for the case of the fixed scalars we have determined the Hawking rate upto a normalization constant. This constant can be determined by the same methods used for the case of the minimal scalars discussed in Chapter 4. We compare the description of the Higgs branch of the effective $U(1)$ gauge theory describing the splitting of (Q_1, Q_5) system into subsystems in terms of Coulomb variables [11] with the supergravity description of the splitting. We find that in both cases the splitting is described by a linear dilaton theory with the same background charge.

In Chapter 4, we finally address the problem of Hawking radiation. We discuss the Hawking radiation of various scalars from the D1/D5 black hole. The near horizon geometry of the D1/D5 black hole is $BTZ \times S^3 \times T^4$ [12] where BTZ refers to a black hole in three-dimensional Anti-de Sitter space discovered by [22]. The mass M and the angular momentum J of the BTZ black hole is given by

$$\begin{aligned} \frac{M}{2} &= L_0 + \bar{L}_0 = \frac{N_R + N_L}{Q_1 Q_5} \\ \frac{J}{2R} &= L_0 - \bar{L}_0 = \frac{N_L - N_R}{Q_1 Q_5} \end{aligned} \tag{1.28}$$

where $R = 2\pi\alpha'(g_s^2 Q_1 Q_5 / V_4)^{1/4}$, N_L and N_R are as defined in (1.25). The metric of the zero mass BTZ black hole is almost identical to that of AdS_3 except for the identification of x_5 with $x_5 + 2\pi R_5$. This difference leads to killing spinors which are periodic in the x_5 coordinate for the BTZ black hole and anti-periodic for the AdS_3 . Thus the SCFT dual to the BTZ is the Ramond sector of the SCFT of the D1/D5 system whereas AdS_3 corresponds to the Neveu-Schwarz sector. The microscopic degrees of freedom for the D1/D5 extremal black hole are the states with $L_0 = N$ and $\bar{L}_0 = 0$ in the Ramond sector of this SCFT. The number of such states agree precisely with the entropy formula (1.13). We argue that the identification of the operator $\mathcal{O}$ corresponding to the minimal and fixed scalars carried out in the Neveu-Schwarz sector remains valid in the Ramond sector of the SCFT. Only non-extremal black holes can Hawking radiate. The general D1/D5 black hole corresponds to a SCFT with $L_0 \neq 0$ and $\bar{L}_0 \neq 0$ over the Ramond vacuum.

We calculate Hawking radiation of minimal scalars from the D1/D5 black hole using the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$. This is done by relating the absorption cross-section of a quanta to the thermal Green's function of the SCFT operator $\mathcal{O}$ which couples to the minimal scalars. The thermal Green's function is determined upto a normalization constant by the conformal dimensions $(h, \bar{h})$ of the operator $\mathcal{O}$ corresponding to the minimal scalars. This method of symmetries makes it possible to derive Hawking radiation of the minimal scalars corresponding to the blow up modes of the orbifold $\mathcal{M}$. In the long string model there is no way of understanding the Hawking radiation of these minimal scalars. This is because these operators are not present in the action of the long string model. The method of using thermal Green's function of the operator $\mathcal{O}$ also makes it clear that upto a normalization constant one can understand Hawking radiation using symmetries.

The precise matching of the Hawking rate computed from the microscopic theory and that computed from supergravity occurs only on fixing the strength of the coupling μ in (1.26). Since the microscopic theory is an effective theory, the precise value of the coupling μ is difficult to fix. We have no first principle derivation of the strength of the coupling. (Note that even in the pion-nucleon model the strength of the coupling g_{eff} could not be fixed by symmetries, one had to appeal to QCD, the microscopic theory relevant for the pion-nucleon model.) We show that consistency with the AdS/CFT correspondence fixes the strength so that the Hawking rate computed from the SCFT based on the orbifold $\mathcal{M}$ and supergravity match exactly. A first principle derivation of the strength of this coupling from the microscopic theory of the D1/D5 system is still an open problem. To this extent a complete understanding of Hawking radiation is still elusive.

We also investigate the Hawking radiation of minimal scalars for the D1/D5 black hole with moduli turned on. We argue that the entropy of the D1/D5 black holes is independent of the moduli. We show that both in the supergravity and in the SCFT the moduli does not modify the Hawking rate [18]. Recently the supergravity solutions of the D1/D5 black hole with moduli were constructed [23]. It would be interesting to verify this prediction for Hawking rate of minimal scalars using the explicit supergravity solution.

It is clear that the method used in this thesis relies only on the symmetry properties

of the SCFT and the near horizon supergravity. The effective SCFT captures the essential symmetries required to reproduce the supergravity result. So it is not a surprise that the SCFT calculations of Hawking radiation done in the limit (1.21) should match with the supergravity calculations done in the limit (1.20).

References

1. For a review of black hole physics and a list of relevant references, see “General Relativity,” by Robert M. Wald, The University of Chicago Press, 1984.
2. S. Hawking, “Gravitational collapsed objects of very low mass,” *Monthly Notices Roy. Astron. Soc.* **152** (1971) 75.
3. A. Salam, “Impact of quantum gravity theory on particle physics,” in *Quantum Gravity: an Oxford Symposium*, eds. C.J. Isham, R. Penrose and D.W. Sciama, Oxford University Press (1975).
4. G. ’t Hooft, “The black hole interpretation of string theory,” *Nucl. Phys.* **B335** (1990) 138.
5. C. Holzhey and F. Wilczek, “Black holes as elementary particles,” *Nucl. Phys.* **B335** (199) 138, hep-th/9202014.
6. L. Susskind, “Some speculations about black hole entropy in string theory,” hep-th/9309145.
7. A. Sen, “Extremal black holes and elementary string states,” *Mod. Phys. Lett.* **A10** (1995) 2081, hep-th/9504147.
8. J. Polchinski, “Dirichlet branes and Ramond-Ramond charges,” *Phys. Rev. Lett.* **75** (1995) 4724, hep-th/9510017.
9. A. Strominger and C. Vafa, “Microscopic origin of the Bekenstein-Hawking entropy,” *Phys. Lett.* **B379** (1996) 99, hep-th/9601029.

10. C. G. Callan and J. Maldacena, “D-brane approach to black hole quantum mechanics,” Nucl. Phys. **B472** (1996) 591, hep-th/9602043.
11. O. Aharony and M. Berkooz, “IR dynamics of $d = 2, \mathcal{N} = (4, 4)$ gauge theories and DLCQ of ‘Little String Theories’,” hep-th/9909101.
12. J. Maldacena and A. Strominger, “Black hole greybody factors and D-brane spectroscopy,” Phys. Rev. **D55** (1997) 861, hep-th/9609026.
13. J. Maldacena and L. Susskind, “D-branes and fat black holes,” Nucl. Phys. **B475** (1996) 679, hep-th/9604042.
14. A. Dhar, G. Mandal and S.R. Wadia, “Absorption vs decay of black holes in string theory and T-symmetry,” Phys. Lett. **B388** (1996) 51, hep-th/9605234.
15. S. R. Das, and S.D. Mathur, “Comparing decay rates for black holes and D-branes,” Nucl. Phys. **B478** (1996) 561, hep-th/9606185; “Interactions involving D-branes,” Nucl. Phys. **B482** (1996) 153, hep-th/9607149.
16. C. G. Callan, S.S. Gubser, I.R. Klebanov and A. A. Tseytlin, “Absorption of fixed scalars and the D-brane approach to black holes,” Nucl. Phys. **B489** (1997) 65, hep-th/9610172.
17. S. F. Hassan and S.R. Wadia, “Gauge theory description of D-brane black holes: emergence of the effective SCFT and Hawking radiation,” Nucl. Phys. **B526** (1998) 311, hep-th/9712213.
18. J.R. David, G. Mandal and S.R. Wadia, “D1/D5 moduli in SCFT and gauge theory and Hawking radiation,” hep-th/9907075.
19. J. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2** (1997) 231, hep-th/9711200.
20. J. de Boer, “Six-dimensional supergravity on $S^3 \times AdS_3$ and 2d conformal field theory,” hep-th/9806104.

- 21. J.R. David, G. Mandal and S.R. Wadia, “Absorption and Hawking radiation of minimal and fixed scalars, and AdS/CFT correspondence,” Nucl. Phys. **B544** (1999) 590, hep-th/9808168.
- 22. M. Banados, C. Teitelboim and J. Zanelli, “Black hole in Three-dimensional space-time,” Phys. Rev. Lett **69** (1992) 1849.
- 23. A. Dhar, G. Mandal, S.R. Wadia and K.P. Yogendran, “D1/D5 system with B-field, noncommutative geometry and the CFT of the Higgs branch,” hep-th/9910194.

Chapter 2

The Microscopic Theory of the D1/D5 system

In this chapter, we will focus on the microscopic description of the D1/D5 system. We will describe the general D1/D5 system with moduli in terms of the $\mathcal{N} = (4, 4)$ SCFT on the resolved orbifold $\mathcal{M}$. We construct all the marginal operators of this SCFT including the blow up modes. We will provide an explicit construction of all short multiplets of single particle states of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$. We also derive an effective $U(1)$ gauge theory describing the dynamics of the splitting of the D1/D5 system into subsystem [1]. We start by reviewing the gauge theory of the D1/D5 system in detail.

2.1 The gauge theory of the D1/D5 system

The gauge theory relevant for understanding the low energy degrees of freedom of the D1/D5 system consists of the massless open string excitations attached to the various D-branes [2, 3]. Consider the Q_1 D1-branes and the Q_5 D5-branes of type IIB string theory in ten-dimensions arranged as follows. The D5-branes are extended along the directions x_5, x_6, x_7, x_8, x_9 and the D1-branes are aligned along the x_5 direction. Let us call the open strings joining the D1-branes among themselves as $(1, 1)$ strings and the strings joining the D5-branes among themselves as $(5, 5)$ strings. The strings joining the D1-branes and the D5-branes are called $(1, 5)$ strings or $(5, 1)$ strings according to their orientations. The boundary conditions of the various open strings at both ends along the various co-ordinates can be read off from the

following table.

co-ordinate	x_0	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_8	x_9
Q_1 D1-branes	N	D	D	D	D	N	D	D	D	D
Q_5 D5-branes	N	D	D	D	D	N	N	N	N	N

(2.1)

Here D stands for Dirichlet boundary conditions and N stands for Neumann boundary conditions. The directions x_6, x_7, x_8, x_9 are compactified on a torus T^4 .

To find the low energy degrees of freedom let us first look at the symmetries preserved by this configuration of D-branes. The $SO(1, 9)$ symmetry of ten-dimensions is broken down to $SO(1, 1) \times SO(4)_E \times SO(4)_I$ by the boundary conditions. The $SO(4)_E$ stands for rotations of the 1, 2, 3, 4 directions and the $SO(4)_I$ stands for rotations of the 6, 7, 8, 9 directions. As the 6, 7, 8, 9 directions are compactified on the torus T^4 , the $SO(4)_I$ symmetry is also broken. But we can still use the $SO(4)_I$ algebra to classify states and organize fields. This configuration preserves eight supersymmetries out of the 32 supersymmetries of the type IIB string theory. Therefore the massless excitations of the various open strings can be organized into fields of $N = 2$ supersymmetry in four-dimensions. Since the size of the T^4 is of the order of string length we can ignore the massive Kaluza-Klein modes on the T^4 . Therefore the low energy effective theory of the D1/D5 system is a $1 + 1$ field theory with eight supersymmetries. We will discuss the field content of this theory as obtained from the various strings joining the D-branes.

$(1, 1)$ strings

The fields corresponding to the massless excitations of the $(1, 1)$ strings can be obtained from the dimensional reduction of $U(Q_1)$ $N = 1$ super Yang-Mills in ten-dimensions to two-dimensions. The bosonic fields of this theory can be organized into the vector multiplet and the hypermultiplet of $N = 2$ theory in four-dimensions as

$$\text{Vector multiplet: } A_0^{(1)}, A_5^{(1)}, \phi_2^{(1)}, \phi_3^{(1)}, \phi_4^{(1)} \quad (2.2)$$

$$\text{Hypermultiplet: } Y_6^{(1)}, Y_7^{(1)}, Y_8^{(1)}, Y_9^{(1)}$$

The $A_0^{(1)}, A_5^{(1)}$ are the $U(Q_1)$ gauge fields in the non-compact directions. The $Y^{(1)}$'s and $\phi^{(1)}$'s are gauge fields in the compact directions of the $N = 1$ super Yang-Mills in ten-dimensions.

They are hermitian $Q_1 \times Q_1$ matrices transforming as adjoints of $U(Q_1)$. The hypermultiplet of $N = 2$ supersymmetry are doublets of the $SU(2)_R$ symmetry of the theory. The $Y^{(1)}$'s can be arranged as doublets under $SU(2)_R$ as

$$N_{a\bar{a}}^{(1)} = \begin{pmatrix} N_{1(a\bar{a})}^{(1)} \\ N_{2(a\bar{a})}^{(1)\dagger} \end{pmatrix} = \begin{pmatrix} Y_{9(a\bar{a})}^{(1)} + iY_{8(a\bar{a})}^{(1)} \\ Y_{7(a\bar{a})}^{(1)} - iY_{6(a\bar{a})}^{(1)} \end{pmatrix} \quad (2.3)$$

where $a, \bar{a}$ runs from $1, \dots, Q_1$.

(5, 5) strings

The field content of these massless open strings is similar to the the (1, 1) strings except for the fact that the gauge group is $U(Q_5)$ instead of $U(Q_1)$. Normally one would have expected the gauge theory of the (5, 5) strings to be a dimensional reduction of $N = 1$ $U(Q_5)$ super Yang-Mills to $5 + 1$ dimensions. Since we are ignoring the Kaluza-Klein modes on T^4 this is effectively a theory in $1 + 1$ dimensions. The vector multiplets and the hypermultiplets are given by

$$\text{Vector multiplet: } A_0^{(5)}, A_5^{(5)}, \phi_2^{(5)}, \phi_3^{(5)}, \phi_4^{(5)} \quad (2.4)$$

$$\text{Hypermultiplet: } Y_6^{(5)}, Y_7^{(5)}, Y_8^{(5)}, Y_9^{(5)}$$

The $A_0^{(5)}, A_5^{(5)}$ are the $U(Q_5)$ gauge fields in the non-compact directions. The $Y^{(5)}$'s and $\phi^{(5)}$'s are gauge fields in the compact directions of the $N = 1$ super Yang-Mills in ten-dimensions. They are hermitian $Q_5 \times Q_5$ matrices transforming as adjoints of $U(Q_5)$. The hypermultiplets $Y^{(5)}$'s can be arranged as doublets under $SU(2)_R$ as

$$N_{b\bar{b}}^{(5)} = \begin{pmatrix} N_{1(b\bar{b})}^{(5)} \\ N_{2(b\bar{b})}^{(5)\dagger} \end{pmatrix} = \begin{pmatrix} Y_{9(b\bar{b})}^{(5)} + iY_{8(b\bar{b})}^{(5)} \\ Y_{7(b\bar{b})}^{(5)} - iY_{6(b\bar{b})}^{(5)} \end{pmatrix} \quad (2.5)$$

where $b, \bar{b}$ runs from $1, \dots, Q_5$.

(1, 5) and (5, 1) strings

From (2.1) we find that these strings have ND boundary conditions along 6, 7, 8, 9 directions. These boundary conditions imply that the world sheet bosons x^6, x^7, x^8, x^9 of the open string have anti-periodic boundary conditions and the world sheet fermions $\psi^6, \psi^7, \psi^8, \psi^9$ are

periodic in the Neveu-Schwarz sector. It can be shown that the net zero point energy of the Neveu-Schwarz sector vanishes. Thus the massless mode is a boson transforming as a spinor of $SO(4)_I$. This gives 2^2 number of bosons. The GSO projection projects out half of these which reduces the number of bosons to 2. The two bosons of the $(1, 5)$ strings and the $(5, 1)$ strings combine to form a complex doublet transforming under the diagonal $SU(2)$ of the $SO(4)_I$. As the hypermultiplets of $N = 2$ theory transform as doublets under $SU(2)_R$, the diagonal $SU(2)$ of $SO(4)_I$ can be identified with the $SU(2)_R$ of the gauge theory. The Chan-Paton factors show that they transform as bi-fundamentals of $U(Q_1)$ and $\overline{U(Q_5)}$. We arrange these hypermultiplets as doublets of the $SU(2)_R$ symmetry of the theory as

$$\chi_{a\bar{b}} = \begin{pmatrix} A_{a\bar{b}} \\ B_{a\bar{b}}^\dagger \end{pmatrix} \quad (2.6)$$

As an aside we note that the fermionic superpartners of these hypermultiplets which arise from the Ramond sector of the massless excitations of $(1, 5)$ and $(5, 1)$ strings carry spinorial indices under $SO(4)_E$ and they are singlets under $SO(4)_I$.

Therefore the gauge theory of the D1/D5 system is a 1+1 dimensional $(4, 4)$ supersymmetric gauge theory with gauge group $U(Q_1) \times U(Q_5)$. The matter content of this theory consists of hypermultiplets $Y^{(1)}$'s, $Y^{(5)}$'s transforming as adjoints of $U(Q_1)$ and $U(Q_5)$ respectively. It also has the hypermultiplets χ 's which transform as bi-fundamentals of $U(Q_1) \times \overline{U(Q_5)}$.

We will now show that this gauge theory has the required degrees of freedom to describe the entropy of the extremal D1/D5 black hole. The D1/D5 bound state is described by the Higgs branch of this gauge theory. The Higgs branch is obtained by giving expectation values to the hypers. This makes the vector multiplets massive. For a supersymmetric vacuum the hypers take values over the surface which is given by setting the superpotential of the gauge theory to zero. Setting the superpotential to zero imposes two sets of D-flatness conditions corresponding to each of the gauge groups $U(Q_1)$ and $U(Q_5)$. The D-terms for the gauge group $U(Q_1)$ is given by

$$\begin{aligned} A_{a\bar{b}} A_{a'\bar{b}}^* - B_{b\bar{a}'} B_{b\bar{a}}^* + [N_1^{(1)}, N_1^{(1)\dagger}]_{a\bar{a}'} - [N_2^{(1)}, N_2^{(1)\dagger}]_{a\bar{a}'} &= 0 \\ A_{a\bar{b}} B_{b\bar{a}'} + [N_1^{(1)}, N_2^{(1)}]_{a\bar{a}'} &= 0 \end{aligned} \quad (2.7)$$

While the D-terms of the gauge group $U(Q_5)$ are

$$\begin{aligned} A_{a\bar{b}'}A_{a\bar{b}}^* - B_{b\bar{a}}B_{b'\bar{a}}^* + [N_1^{(5)}, N_1^{(5)\dagger}]_{b\bar{b}'} - [N_2^{(5)}, N_2^{(5)\dagger}]_{b\bar{b}'} &= 0 \\ A_{a\bar{b}'}B_{b\bar{a}} + [N_1^{(5)}, N_2^{(5)}]_{b\bar{b}'} &= 0 \end{aligned} \quad (2.8)$$

where a' runs from $1, \dots, Q_1$ and b' runs from $1, \dots, Q_5$.

The total number of bosonic degrees of freedom from all the hypermultiplets is

$$4Q_1^2 + 4Q_5^2 + 4Q_1Q_5 \quad (2.9)$$

The first equation in (2.7) is real and the second equation in (2.7) is complex. The total number of constraints imposed by (2.7) is $3Q_1^2$. Similarly the set of D-term equations in (2.8) imposes $3Q_5^2$ constraints. Equations (2.7) and (2.8) have the same trace parts corresponding to the vanishing of $U(1)$ D-terms, namely,

$$\begin{aligned} A_{a\bar{b}}A_{a\bar{b}}^* - B_{b\bar{a}}B_{b\bar{a}}^* &= 0 \\ A_{a\bar{b}}B_{a\bar{b}} &= 0 \end{aligned} \quad (2.10)$$

which are three real equations. Therefore, the vanishing of D-terms imposes $3Q_1^2 + 3Q_5^2 - 3$ constraints on the fields. One can use the gauge symmetry $U(Q_1)$ and $U(Q_5)$ to remove another $Q_1^2 + Q_5^2 - 1$ degrees of freedom. The number of degrees of freedom fixed by the gauge symmetry is less than the number of generators of $U(Q_1)$ and $U(Q_5)$ by one. This is because all the hypermultiplets are invariant under the diagonal $U(1)$ of $U(Q_1) \times U(Q_5)$. After gauge fixing, the number of gauge invariant bosonic degrees of freedom to parameterize the moduli space is $4(Q_1Q_5 + 1)$.

We are interested in low energy black hole processes so it is sufficient to study the SCFT of the Higgs branch. The SCFT will have $\mathcal{N} = (4, 4)$ SUSY with central charge $6(Q_1Q_5 + 1)$ on a target space $\mathcal{M}$. To find the microstates corresponding to the extremal D1/D5 black hole we look for states with $L_0 = N$ and $\bar{L}_0 = 0$. The asymptotic number of distinct states of this SCFT given by Cardy's formula

$$\Omega = \exp(2\pi\sqrt{Q_1Q_5N}) \quad (2.11)$$

From the Boltzmann formula one obtains

$$S = 2\pi\sqrt{Q_1 Q_5 N} \quad (2.12)$$

This exactly reproduces the Bekenstein-Hawking entropy (1.13) of the extremal D1/D5 black hole.

2.2 The instanton moduli space

In the previous section we found that the Higgs branch of the gauge theory of the D1/D5 system flows in the infrared to $\mathcal{N} = (4, 4)$ SCFT on a target space $\mathcal{M}$ with central charge $6Q_1 Q_5$. For black hole processes like Hawking radiation it is important to know the target space $\mathcal{M}$. In this section we review the arguments which show that the target space $\mathcal{M}$ is a resolution of the orbifold $(\tilde{T}^4)^{Q_1 Q_5} / S(Q_1 Q_5)$. ($\tilde{T}^4$ can be different from the compactification torus T^4 .)

The Q_1 D1-branes can be thought of as Q_1 instantons in the $5 + 1$ dimensional $U(Q_5)$ super Yang-Mills theory of the Q_5 D5-branes [4]. To see this note that the DBI action of the D5-branes have a coupling

$$\int d^6 x \ C^{(2)} \wedge \text{Tr}[F^{(5)} \wedge F^{(5)}] \quad (2.13)$$

The non-trivial gauge configurations which are independent of x^0, x^5 and have zero values of $A_0^{(5)}$ and $A_5^{(5)}$ but non-zero values of $\text{Tr}[F^{(5)} \wedge F^{(5)}]$ act as sources of the Ramond-Ramond two-form $C_{05}^{(2)}$. If these gauge field configurations have to preserve half the supersymmetries of the D5-brane action they should be self dual. Thus they are instanton solutions of four-dimensional Euclidean Yang-Mills of the 6, 7, 8, 9 directions.

Additional evidence for this comes from the fact that the integral property of $\text{Tr}[F^{(5)} \wedge F^{(5)}]$ corresponds to the quantization of the D1-branes charge. The action for a Q_1 instanton solution is Q_1/g_{YM}^2 . This agrees with the tension of Q_1 D1-branes, namely Q_1/g_s . If one is dealing with non-compact D5-branes and D1-branes it is seen that the D-flatness conditions of the D1-brane theory is identical to the ADHM construction of Q_1 instantons of $U(Q_5)$ gauge theory [5].

From the discussion in the preceding paragraphs we conclude that $\mathcal{M}$ the target space of SCFT can be thought of as the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on T^4 . This moduli space is known to be the Hilbert scheme of $Q_1 Q_5$ points on $\tilde{T}^4$ [6]. $\tilde{T}^4$ can be different from the compactification torus T^4 . This is a smooth resolution of the singular orbifold $(\tilde{T}^4)^{Q_1 Q_5}/S(Q_1 Q_5)$. We will provide physically motivated evidence for the fact that the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on T^4 is a smooth resolution of the orbifold $(\tilde{T}^4)^{Q_1 Q_5}/S(Q_1 Q_5)$ using string dualities. The evidence is topological and it comes from realising that the cohomology of $\mathcal{M}$ is the degeneracy of the ground states of the D1/D5 gauge theory. We can calculate this degeneracy in two ways. One is by explicitly counting the cohomology of $(\tilde{T}^4)^{Q_1 Q_5}/S(Q_1 Q_5)$ [7]. The second method is to use string dualities as discussed below. Both these methods give identical answers. Thus at least at the level of cohomology we are able to verify that the moduli space is $(\tilde{T}^4)^{Q_1 Q_5}/S(Q_1 Q_5)$.

Consider type IIB string theory compactified on $S^1 \times T^4$ with a fundamental string having Q_5 units of winding along x^6 and Q_1 units of momentum along x^6 . On performing the sequence of dualities $ST_{6789}ST_{56}$ we can map the fundamental string to the D1/D5 system (we can de-compactify the x^5 direction finally) with Q_1 D1-branes along x^5 and Q_5 D5-branes along x^5, x^6, x^7, x^8, x^9 [8]. Therefore using this U-duality sequence the BPS states of this fundamental string (that is, states with either purely left moving or right moving oscillators) maps to ground states of the D1/D5 system. The number of ground states of the D1/D5 system is given by the dimension of the cohomology of $\mathcal{M}$. From the perturbative string degeneracy counting the generating function of BPS states with left moving oscillator number N_L is given by

$$\sum_{N_L=0}^{\infty} d(N_L) q^{N_L} = 256 \times \prod_{n=1}^{\infty} \left(\frac{1+q^n}{1-q^n} \right)^8 \quad (2.14)$$

where $d(N_L)$ refers to the degeneracy of states with left moving oscillator number N_L . The D1/D5 system is U-dual to the perturbative string with $N_L = Q_1 Q_5$.

Explicit counting of the cohomology of $\mathcal{M}$ gives $d(Q_1 Q_5)/256$. The factor 256 comes from quantization of the center of mass coordinate along 1, 2, 3, 4 directions and the 6, 7, 8, 9 directions. The center of mass coordinate is represented by the relative $U(1)$ of $U(Q_1) \times U(Q_5)$. Therefore the low energy theory of the bound D1/D5 system is a SCFT on the target

space

$$R^4 \times T^4 \times (\tilde{T}^4)^{Q_1 Q_5} / S(Q_1 Q_5) \quad (2.15)$$

From now on we will suppress the center of mass coordinate and $\mathcal{M}$ will denote

$$(\tilde{T}^4)^{Q_1 Q_5} / S(Q_1 Q_5) \quad (2.16)$$

2.3 The SCFT on the orbifold $\mathcal{M}$

The $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ is described by the free Lagrangian

$$S = \frac{1}{2} \int d^2 z \left[\partial x_A^i \bar{\partial} x_{i,A} + \psi_A^i(z) \bar{\partial} \psi_A^i(z) + \tilde{\psi}_A^i(\bar{z}) \partial \tilde{\psi}_A^i(\bar{z}) \right] \quad (2.17)$$

Here i runs over the $\tilde{T}^4$ coordinates 1,2,3,4 and $A = 1, 2, \dots, Q_1 Q_5$ labels various copies of the four-torus. The symmetric group $S(Q_1 Q_5)$ acts by permuting the copy indices. It introduces various twisted sectors which we will discuss later. The free field realization of this SCFT has $\mathcal{N} = (4, 4)$ superconformal symmetry. To set up our notations and conventions we review the $\mathcal{N} = 4$ superconformal algebra.

2.3.1 The $\mathcal{N} = 4$ superconformal algebra

The algebra is generated by the stress energy tensor, four supersymmetry currents, and a local $SU(2)$ R symmetry current. The operator product expansions(OPE) of the algebra with central charge c are given by (See for example [9].)

$$\begin{aligned} T(z)T(w) &= \frac{\partial T(w)}{z-w} + \frac{2T(w)}{(z-w)^2} + \frac{c}{2(z-w)^4}, \\ G^a(z)G^{b\dagger}(w) &= \frac{2T(w)\delta_{ab}}{z-w} + \frac{2\bar{\sigma}_{ab}^i \partial J^i}{z-w} + \frac{4\bar{\sigma}_{ab}^i J^i}{(z-w)^2} + \frac{2c\delta_{ab}}{3(z-w)^3}, \\ J^i(z)J^j(w) &= \frac{i\epsilon^{ijk} J^k}{z-w} + \frac{c}{12(z-w)^2}, \\ T(z)G^a(w) &= \frac{\partial G^a(w)}{z-w} + \frac{3G^a(z)}{2(z-w)^2}, \\ T(z)G^{a\dagger}(w) &= \frac{\partial G^{a\dagger}(w)}{z-w} + \frac{3G^{a\dagger}(z)}{2(z-w)^2}, \\ T(z)J^i(w) &= \frac{\partial J^i(w)}{z-w} + \frac{J^i}{(z-w)^2}, \end{aligned} \quad (2.18)$$

$$\begin{aligned}
J^i(z)G^a(w) &= \frac{G^b(z)(\sigma^i)^{ba}}{2(z-w)}, \\
J^i(z)G^{a\dagger}(w) &= -\frac{(\sigma^i)^{ab}G^{b\dagger}(w)}{2(z-w)}
\end{aligned}$$

Here $T(z)$ is the stress energy tensor, $G^a(z), G^{b\dagger}(z)$ the $SU(2)$ doublet of supersymmetry generators and $J^i(z)$ the $SU(2)$ R symmetry current. The σ 's stand for Pauli matrices and the $\bar{\sigma}$'s stand for the complex conjugates of Pauli matrices. In the free field realization described below, the above holomorphic currents occur together with their antiholomorphic counterparts, which we will denote by $\tilde{J}(\bar{z}), \tilde{G}(\bar{z})$ and $\tilde{T}(\bar{z})$. In particular, the R-parity group will be denoted by $SU(2)_R \times \widetilde{SU(2)}_R$.

2.3.2 Free field realization of $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$

A free field realization of the $\mathcal{N} = 4$ superconformal algebra with $c = 6Q_1Q_5$ can be constructed out of Q_1Q_5 copies of four real fermions and bosons. The generators are given by

$$\begin{aligned}
T(z) &= \partial X_A(z)\partial X_A^\dagger(z) + \frac{1}{2}\Psi_A(z)\partial\Psi_A^\dagger(z) - \frac{1}{2}\partial\Psi_A(z)\Psi_A^\dagger(z) \\
G^a(z) &= \begin{pmatrix} G^1(z) \\ G^2(z) \end{pmatrix} = \sqrt{2}\begin{pmatrix} \Psi_A^1(z) \\ \Psi_A^2(z) \end{pmatrix} \partial X_A^2(z) + \sqrt{2}\begin{pmatrix} -\Psi_A^{2\dagger}(z) \\ \Psi_A^{1\dagger}(z) \end{pmatrix} \partial X_A^1(z) \\
J_R^i(z) &= \frac{1}{2}\Psi_A(z)\sigma^i\Psi_A^\dagger(z)
\end{aligned} \tag{2.19}$$

We will use the following notation for the zero mode of the R-parity current:

$$J_R^i = \frac{1}{2} \int \frac{dz}{2\pi i} \Psi_A(z) \sigma^i \Psi_A^\dagger(z) \tag{2.20}$$

In the above the summation over A which runs from 1 to Q_1Q_5 is implied. The bosons X and the fermions Ψ are

$$\begin{aligned}
X_A(z) &= (X_A^1(z), X_A^2(z)) = \sqrt{1/2}(x_A^1(z) + ix_A^2(z), x_A^3(z) + ix_A^4(z)), \\
\Psi_A(z) &= (\Psi_A^1(z), \Psi_A^2(z)) = \sqrt{1/2}(\psi_A^1(z) + i\psi_A^2(z), \psi_A^3(z) + i\psi_A^4(z)) \\
X_A^\dagger(z) &= \begin{pmatrix} X_A^{1\dagger}(z) \\ X_A^{2\dagger}(z) \end{pmatrix} = \sqrt{\frac{1}{2}} \begin{pmatrix} x_A^1(z) - ix_A^2(z) \\ x_A^2(z) - ix_A^3(z) \end{pmatrix} \\
\Psi_A^\dagger(z) &= \begin{pmatrix} \Psi_A^{1\dagger}(z) \\ \Psi_A^{2\dagger}(z) \end{pmatrix} = \sqrt{\frac{1}{2}} \begin{pmatrix} \psi_A^1(z) - i\psi_A^2(z) \\ \psi_A^3(z) - i\psi_A^4(z) \end{pmatrix}
\end{aligned} \tag{2.21}$$

2.3.3 The $SO(4)$ algebra

In addition to the local R symmetry the free field realization of the $\mathcal{N} = 4$ superconformal algebra has additional global symmetries which can be used to classify the states. There are 2 global $SU(2)$ symmetries which correspond to the $SO(4)$ rotations of the 4 bosons x^i . The corresponding charges are given by

$$\begin{aligned} I_1^i &= \frac{1}{4} \int \frac{dz}{2\pi i} X_A \sigma^i \partial X_A^\dagger - \frac{1}{4} \int \frac{dz}{2\pi i} \partial X_A \sigma^i X_A^\dagger + \frac{1}{2} \int \frac{dz}{2\pi i} \Phi_A \sigma^i \Phi_A^\dagger \\ I_2^i &= \frac{1}{4} \int \frac{dz}{2\pi i} \mathcal{X}_A \sigma^i \partial \mathcal{X}_A^\dagger - \frac{1}{4} \int \frac{dz}{2\pi i} \partial \mathcal{X}_A \sigma^i \mathcal{X}_A^\dagger \end{aligned} \quad (2.22)$$

Here

$$\begin{aligned} \mathcal{X}_A &= (X_A^1, -X_A^{2\dagger}) & \mathcal{X}^\dagger &= \begin{pmatrix} X_A^{1\dagger} \\ -X_A^2 \end{pmatrix} \\ \Phi_A &= (\Psi_A^1, \Psi_A^{2\dagger}) & \Phi_A^\dagger &= \begin{pmatrix} \Psi_A^{1\dagger} \\ \Psi_A^2 \end{pmatrix}. \end{aligned} \quad (2.23)$$

These charges are generators of $SU(2) \times SU(2)$ algebra:

$$\begin{aligned} [I_1^i, I_1^j] &= i\epsilon^{ijk} I_1^k & [I_2^i, I_2^j] &= i\epsilon^{ijk} I_2^k \\ [I_1^i, I_2^j] &= 0 \end{aligned} \quad (2.24)$$

The commutation relation of these new global charges with the various local charges are given below

$$\begin{aligned} [I_1^i, G^a(z)] &= 0 & [I_1^i, G^{a\dagger}(z)] &= 0 \\ [I_1^i, T(z)] &= 0 & [I_1^i, J(z)] &= 0 \\ [I_2^i, \mathcal{G}^a(z)] &= \frac{1}{2} \mathcal{G}^b(z) \sigma_{ba}^i & [I_2^i, \mathcal{G}^{a\dagger}(z)] &= -\frac{1}{2} \sigma_{ab}^i \mathcal{G}^{b\dagger}(z) \\ [I_2^i, T(z)] &= 0 & [I_2^i, J(z)] &= 0 \end{aligned} \quad (2.25)$$

where

$$\mathcal{G} = (G^1, G^{2\dagger}) \quad \mathcal{G}^\dagger = \begin{pmatrix} G^{1\dagger} \\ G^2 \end{pmatrix} \quad (2.26)$$

The following commutations relation show that the bosons transform as $(\mathbf{2}, \mathbf{2})$ under $SU(2)_{I_1} \times SU(2)_{I_2}$

$$\begin{aligned} [I_1^i, X_A^a] &= \frac{1}{2} X_A^b \sigma_{ba}^i & [I_1^i, X_A^{a\dagger}] &= -\frac{1}{2} \sigma_{ab}^i X_A^{b\dagger} \\ [I_2^i, \mathcal{X}_A^a] &= \frac{1}{2} \mathcal{X}_A^b \sigma_{ba}^i & [I_2^i, \mathcal{X}_A^{a\dagger}] &= -\frac{1}{2} \sigma_{ab}^i \mathcal{X}_A^{b\dagger} \end{aligned} \quad (2.27)$$

The fermions transform as $(\mathbf{2}, \mathbf{1})$ under $SU(2)_{I_1} \times SU(2)_{I_2}$ as can be seen from the commutation relations given below.

$$\begin{aligned} [I_1^i, \Phi_A^a] &= \frac{1}{2} \Phi_A^b \sigma_{ba}^i & [I_1^i, \Phi_A^{a\dagger}] &= -\frac{1}{2} \sigma_{ab}^i \Phi_A^{b\dagger} \\ [I_2^i, \Psi^a] &= 0 & [I_2^i, \bar{\Psi}^a] &= 0 \end{aligned} \quad (2.28)$$

We are interested in studying the states of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$. The classification of the states and their symmetry properties can be analyzed by studying the states of a free field realization of a $\mathcal{N} = (4, 4)$ SCFT on $R^{4Q_1Q_5}/S(Q_1Q_5)$. This is realized by considering the holomorphic and the anti-holomorphic $\mathcal{N} = 4$ superconformal algebra with $c = \bar{c} = 6Q_1Q_5$ constructed out of Q_1Q_5 copies of four real fermions and bosons. So we have an anti-holomorphic component for each field, generator and charges discussed above. These are labelled by the same symbols used for the holomorphic components but distinguished by a tilde.

The charges I_1, I_2 constructed above generate $SO(4)$ transformations only on the *holomorphic* bosons $X_A(z)$. Similarly, we can construct charges $\tilde{I}_1, \tilde{I}_2$ which generate $SO(4)$ transformations only on the *antiholomorphic* bosons $\widetilde{X}_A(\bar{z})$. Normally one would expect these charges to give rise to a global $SO(4)_{hol} \times SO(4)_{antihol}$ symmetry. However, the kinetic term of the bosons in the free field realization is not invariant under independent holomorphic and antiholomorphic $SO(4)$ rotations. It is easy to see, for example by using the Noether procedure, that there is a residual $SO(4)$ symmetry generated by the charges

$$J_I = I_1 + \tilde{I}_1 \quad \tilde{J}_I = I_2 + \tilde{I}_2 \quad (2.29)$$

We will denote this symmetry as $SO(4)_I = SU(2)_I \times \widetilde{SU(2)}_I$, where the $SU(2)$ factors are generated by $J_I, \tilde{J}_I$. These charges satisfy the property that (a) they correspond to $SO(4)$ transformations of the bosons $X_A(z, \bar{z}) = X_A(z) + \widetilde{X}_A(\bar{z})$ and (b) they fall into representations of the $\mathcal{N} = (4, 4)$ algebra (as can be proved by using the commutation relations (2.27) of the I 's). The bosons $X(z, \bar{z})$ transform as $(\mathbf{2}, \mathbf{2})$ under $SU(2)_I \times \widetilde{SU(2)}_I$.

2.3.4 The supergroup $SU(1, 1|2)$

The global part of the $\mathcal{N} = 4$ superconformal algebra forms the supergroup $SU(1, 1|2)$. Let $L_{\pm,0}$, $J_R^{(1),(2),(3)}$ be the global charges of the currents $T(z)$ and $J_R^{(i)}(z)$ and $G_{1/2,-1/2}^a$ the global charges of the supersymmetry currents $G^a(z)$ in the Neveu-Schwarz sector. From the OPE's (2.18) we obtain the following commutation relations for the global charges.

$$\begin{aligned}
[L_0, L_{\pm}] &= \mp L_{\pm} & [L_1, L_{-1}] &= 2L_0 & (2.30) \\
\{G_{1/2}^a, G_{-1/2}^{b\dagger}\} &= 2\delta^{ab}L_0 + 2\sigma_{ab}^i J_R^{(i)} \\
\{G_{-1/2}^a, G_{1/2}^{b\dagger}\} &= 2\delta^{ab}L_0 - 2\sigma_{ab}^i J_R^{(i)} \\
[J_R^{(i)}, J_R^{(j)}] &= i\epsilon^{ijk} J_R^{(k)} \\
[L_0, G_{\pm 1/2}^a] &= \mp \frac{1}{2} G_{\pm 1/2}^a & [L_0, G_{\pm 1/2}^{a\dagger}] &= \mp \frac{1}{2} G_{\pm 1/2}^{a\dagger} \\
[L_+, G_{1/2}^a] &= 0 & [L_-, G_{-1/2}^a] &= 0 \\
[L_-, G_{1/2}^a] &= -G_{-1/2}^a & [L_+, G_{-1/2}^a] &= G_{1/2}^a \\
[L_+, G_{1/2}^{a\dagger}] &= 0 & [L_-, G_{1/2}^{a\dagger}] &= 0 \\
[L_-, G_{1/2}^{a\dagger}] &= -G_{-1/2}^{a\dagger} & [L_+, G_{-1/2}^{a\dagger}] &= G_{1/2}^{a\dagger} \\
[J_R^{(i)}, G_{\pm 1/2}^a] &= \frac{1}{2} G_{\pm 1/2}^b (\sigma^i)^{ba} & [J_R^{(i)}, G_{\pm 1/2}^{a\dagger}] &= -\frac{1}{2} (\sigma^i)^{ba} G_{\pm 1/2}^{b\dagger}
\end{aligned}$$

The above commutation relations form the algebra of the supergroup $SU(1, 1|2)$. The global part of the $\mathcal{N} = (4, 4)$ superconformal algebra form the super group $SU(1, 1|2) \times SU(1, 1|2)$.

2.4 Short multiplets of $SU(1, 1|2)$

The representations of the supergroup $SU(1, 1|2)$ are classified according to the conformal weight and $SU(2)_R$ quantum number. The highest weight states $|\text{hw}\rangle = |h, \mathbf{j}_R, j_R^3 = j_R\rangle$ satisfy the following properties

$$\begin{aligned}
L_1|\text{hw}\rangle &= 0 & L_0|\text{hw}\rangle &= h|\text{hw}\rangle & (2.31) \\
J_R^{(+)}|\text{hw}\rangle &= 0 & J_R^{(3)}|\text{hw}\rangle &= j_R|\text{hw}\rangle \\
G_{1/2}^a|\text{hw}\rangle &= 0 & G_{1/2}^{a\dagger}|\text{hw}\rangle &= 0
\end{aligned}$$

where $J_R^+ = J_R^{(1)} + iJ_R^{(2)}$. Highest weight states which satisfy $G_{-1/2}^{2\dagger}|\text{hw}\rangle = 0$, $G_{-1/2}^1|\text{hw}\rangle = 0$ are chiral primaries. They satisfy $h = j$. We will denote these states as $|\text{hw}\rangle_S$. Short multiplets are generated from the chiral primaries through the action of the raising operators $J_-, G_{-1/2}^{1\dagger}$ and $G_{-1/2}^2$. The structure of the short multiplet is given below

States	j	L_0	Degeneracy
$ \text{hw}\rangle_S$	h	h	$2h + 1$
$G_{-1/2}^{1\dagger} \text{hw}\rangle_S, G_{-1/2}^2 \text{hw}\rangle_S$	$h - 1/2$	$h + 1/2$	$2h + 2h = 4h$
$G_{-1/2}^{1\dagger}G_{-1/2}^2 \text{hw}\rangle_S$	$h - 1$	$h + 1$	$2h - 1$

(2.32)

The short multiplets of the supergroup $SU(1,1|2) \times SU(1,1|2)$ are obtained by the tensor product of the above multiplet. We denote the short multiplet of $SU(1,1|2) \times SU(1,1|2)$ as $(\mathbf{2h} + \mathbf{1}, \mathbf{2h}' + \mathbf{1})_{\mathbf{S}}$. These stand for the degeneracy of the bottom component, the top row in (2.32). The top component of the short multiplet are the states belonging to the last row in (2.32). The short multiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ is special, it terminates at the middle row of (2.32). For this case, the top component is the middle row. These states have $h = \bar{h} = 1$ and transform as $(\mathbf{1}, \mathbf{1})$ of $SU(2)_R \times \widetilde{SU(2)}_R$. There are 4 such states for each $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$.

2.5 The resolutions of the symmetric product

We would like to discuss D1/D5 systems with various moduli turned on. To do this we will construct marginal operators of the $\mathcal{N} = (4, 4)$ SCFT on the symmetric product orbifold $\mathcal{M}$. We will find the four operators which correspond to resolution of the orbifold singularity.

2.5.1 The untwisted sector

Let us first focus on the operators constructed from the untwisted sector. The operators of lowest conformal weight are

$$\begin{aligned}
\Psi_A^1(z) \tilde{\Psi}_A^1(\bar{z}) & \quad \Psi_A^1(z) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \\
\Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) & \quad \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^{2\dagger}(\bar{z})
\end{aligned}
\tag{2.33}$$

where summation over A is implied. These four operators have conformal dimension $(h, \bar{h}) = (1/2, 1/2)$ and $(j_R^3, \tilde{j}_R^3) = (1/2, 1/2)$ under the R-symmetry $SU(2)_R \times \widetilde{SU(2)}_R$. Since $(h, \bar{h}) =$

$(j_R^3, \tilde{j}_R^3)$, these operators are chiral primaries and have non-singular operator product expansions (OPE) with the supersymmetry currents $G^1(z), G^{2\dagger}(z), \tilde{G}^1(\bar{z}), \tilde{G}^{2\dagger}(\bar{z})$. These properties indicate that they belong to the bottom component of the short multiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$. Each of the four chiral primaries gives rise to four top components of the short multiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$. They are given by the leading pole $((z-w)^{-1}(\bar{z}-\bar{w})^{-1})$ in the OPE's

$$\begin{aligned} G^2(z)\tilde{G}^2(\bar{z})\mathcal{P}(w, \bar{w}) & \quad G^2(z)\tilde{G}^{1\dagger}(\bar{z})\mathcal{P}(w, \bar{w}) \\ G^{1\dagger}(z)\tilde{G}^2(\bar{z})\mathcal{P}(w, \bar{w}) & \quad G^{1\dagger}(z)\tilde{G}^{1\dagger}(\bar{z})\mathcal{P}(w, \bar{w}) \end{aligned} \quad (2.34)$$

where $\mathcal{P}$ stands for any of the four chiral primaries in (2.33). From the superconformal algebra it is easily seen that the top components constructed above have weights $(1, 1)$ and transform as $(\mathbf{1}, \mathbf{1})$ under $SU(2)_R \times \widetilde{SU}(2)_R$. The OPE's (2.34) can be easily evaluated. We find that the 16 top components of the $4(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ short multiplets are $\partial x_A^i \bar{\partial} x_A^j$.

We classify the above operators belonging to the top component according to representations of (a) the $SO(4)_I$ rotational symmetry of the $\tilde{T}^4$, (The four torus $\tilde{T}^4$ breaks this symmetry but we assume the target space is R^4 for the classification of states) (b) R symmetry of the SCFT and (c) the conformal weights. As all of these operators belong to the top component of $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ the only property which distinguishes them is the representation under $SO(4)_I$. The quantum numbers of these operators under the various symmetries are

Operator	$SU(2)_I \times \widetilde{SU}(2)_I$	$SU(2)_R \times \widetilde{SU}(2)_R$	$(h, \bar{h})$
$\partial x_A^{\{i}(z)\bar{\partial} x_A^{j\}}(\bar{z}) - \frac{1}{4}\delta^{ij}\partial x_A^k(z)\bar{\partial} x_A^k(\bar{z})$	$(\mathbf{3}, \mathbf{3})$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$
$\partial x_A^{[i}(z)\bar{\partial} x_A^{j]}(\bar{z})$	$(\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \mathbf{3})$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$
$\partial x_A^i(z)\bar{\partial} x_A^i(\bar{z})$	$(\mathbf{1}, \mathbf{1})$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$

(2.35)

Therefore we have 16 marginal operators from the untwisted sector. As these are top components they can be added to the free SCFT as perturbations without violating the $\mathcal{N} = (4, 4)$ supersymmetry.

2.5.2 Z_2 twists.

We now construct the marginal operators from the various twisted sectors of the orbifold SCFT. The twist fields of the SCFT on the orbifold $\mathcal{M}$ are labeled by the conjugacy classes of the symmetric group $S(Q_1 Q_5)$ [10, 11]. The conjugacy classes consist of cyclic groups of

various lengths. The various conjugacy classes and the multiplicity in which they occur in $S(Q_1 Q_5)$ can be found from the solutions of the equation

$$\sum n N_n = Q_1 Q_5 \quad (2.36)$$

where n is the length of the cycle and N_n is the multiplicity of the cycle. Consider the simplest nontrivial conjugacy class which is given by $N_1 = Q_1 Q_5 - 2, N_2 = 1$ and the rest of $N_n = 0$. A representative element of this class is

$$(X_1 \rightarrow X_2, X_2 \rightarrow X_1), X_3 \rightarrow X_3, \dots, X_{Q_1 Q_5} \rightarrow X_{Q_1 Q_5} \quad (2.37)$$

Here the X_A 's are related to the x_A 's appearing in the action (2.17) by (2.21) in.

To exhibit the singularity of this group action we go over to the following new coordinates

$$X_{cm} = X_1 + X_2 \text{ and } \phi = X_1 - X_2 \quad (2.38)$$

Under the group action (2.37) X_{cm} is invariant and $\phi \rightarrow -\phi$. Thus the singularity is *locally* of the type R^4/Z_2 . The bosonic twist operators for this orbifold singularity are given by following OPE's [12]

$$\begin{aligned} \partial\phi^1(z)\sigma^1(w, \bar{w}) &= \frac{\tau^1(w, \bar{w})}{(z-w)^{1/2}} & \partial\phi^{1\dagger}(z)\sigma^1(w, \bar{w}) &= \frac{\tau'^1(w, \bar{w})}{(z-w)^{1/2}} \\ \partial\phi^2(z)\sigma^2(w, \bar{w}) &= \frac{\tau^2(w, \bar{w})}{(z-w)^{1/2}} & \partial\phi^{2\dagger}(z)\sigma^2(w, \bar{w}) &= \frac{\tau'^2(w, \bar{w})}{(z-w)^{1/2}} \\ \bar{\partial}\tilde{\phi}^1(\bar{z})\sigma^1(w, \bar{w}) &= \frac{\tilde{\tau}^1(w, \bar{w})}{(\bar{z}-\bar{w})^{1/2}} & \bar{\partial}\tilde{\phi}^{1\dagger}(\bar{z})\sigma^1(w, \bar{w}) &= \frac{\tilde{\tau}'^1(w, \bar{w})}{(\bar{z}-\bar{w})^{1/2}} \\ \bar{\partial}\tilde{\phi}^2(\bar{z})\sigma^2(w, \bar{w}) &= \frac{\tilde{\tau}^2(w, \bar{w})}{(\bar{z}-\bar{w})^{1/2}} & \bar{\partial}\tilde{\phi}^{2\dagger}(\bar{z})\sigma^2(w, \bar{w}) &= \frac{\tilde{\tau}'^2(w, \bar{w})}{(\bar{z}-\bar{w})^{1/2}} \end{aligned} \quad (2.39)$$

The τ 's are excited twist operators. The fermionic twists are constructed from bosonized currents defined by

$$\begin{aligned} \chi^1(z) &= e^{iH^1(z)} & \chi^{1\dagger}(z) &= e^{-iH^1(z)} \\ \chi^2(z) &= e^{iH^2(z)} & \chi^{2\dagger}(z) &= e^{-iH^2(z)} \end{aligned} \quad (2.40)$$

Where the χ 's, defined as $\Psi_1 - \Psi_2$, are the superpartners of the bosons ϕ .

From the above we construct the supersymmetric twist fields which act both on fermions and bosons as follows:

$$\begin{aligned}
\Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})} &= \sigma^1(z, \bar{z}) \sigma^2(z, \bar{z}) e^{iH^1(z)/2} e^{-iH^2(z)/2} e^{i\tilde{H}^1(\bar{z})/2} e^{-i\tilde{H}^2(\bar{z})/2} \\
\Sigma_{(12)}^{(\frac{1}{2}, -\frac{1}{2})} &= \sigma^1(z, \bar{z}) \sigma^2(z, \bar{z}) e^{iH^1(z)/2} e^{-iH^2(z)/2} e^{-i\tilde{H}^1(\bar{z})/2} e^{i\tilde{H}^2(\bar{z})/2} \\
\Sigma_{(12)}^{(-\frac{1}{2}, \frac{1}{2})} &= \sigma^1(z, \bar{z}) \sigma^2(z, \bar{z}) e^{-iH^1(z)/2} e^{+iH^2(z)/2} e^{i\tilde{H}^1(\bar{z})/2} e^{-i\tilde{H}^2(\bar{z})/2} \\
\Sigma_{(12)}^{(-\frac{1}{2}, -\frac{1}{2})} &= \sigma^1(z, \bar{z}) \sigma^2(z, \bar{z}) e^{-iH^1(z)/2} e^{+iH^2(z)/2} e^{-i\tilde{H}^1(\bar{z})/2} e^{+i\tilde{H}^2(\bar{z})/2}
\end{aligned} \tag{2.41}$$

The subscript (12) refers to the fact that these twist operators were constructed for the representative group element (2.37) which exchanges the 1 and 2 labels of the coordinates of $\tilde{T}^4$. The superscript stands for the $(j_R^3, \tilde{j}_R^3)$ quantum numbers. The twist operators for the orbifold $\mathcal{M}$ belonging to the conjugacy class under consideration is obtained by summing over these Z_2 twist operators for all representative elements of this class.

$$\Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})} = \sum_{i=1}^{Q_1 Q_5} \sum_{j=1, j \neq i}^{Q_1 Q_5} \Sigma_{(ij)}^{(\frac{1}{2}, \frac{1}{2})} \tag{2.42}$$

We can define the rest of the twist operators for the orbifold in a similar manner. The conformal dimensions of these operators are $(1/2, 1/2)$. They transform as $(\mathbf{2}, \mathbf{2})$ under the $SU(2)_R \times \widetilde{SU(2)}_R$ symmetry of the SCFT. They belong to the bottom component of the short multiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{s}}$. The operator $\Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})}$ is a chiral primary. As before the 4 top components of this short multiplet, which we denote by

$$\begin{aligned}
&T^{(\frac{1}{2}, \frac{1}{2})}, \quad T^{(\frac{1}{2}, -\frac{1}{2})} \\
&T^{(-\frac{1}{2}, \frac{1}{2})}, \quad T^{(-\frac{1}{2}, -\frac{1}{2})}
\end{aligned} \tag{2.43}$$

are given by the leading pole in the following OPE's respectively

$$\begin{aligned}
&G^2(z) \tilde{G}^2(\bar{z}) \Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})}(w, \bar{w}), \quad G^2(z) \tilde{G}^{1\dagger}(\bar{z}) \Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})}(w, \bar{w}), \\
&G^{1\dagger}(z) \tilde{G}^2(\bar{z}) \Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})}(w, \bar{w}), \quad G^{1\dagger}(z) \tilde{G}^{1\dagger}(\bar{z}) \Sigma_{(12)}^{(\frac{1}{2}, \frac{1}{2})}(w, \bar{w})
\end{aligned} \tag{2.44}$$

These are the 4 blow up modes of the R^4/Z_2 singularity [14] and they have conformal weight $(1, 1)^1$. They transform as $(\mathbf{1}, \mathbf{1})$ under the $SU(2)_R \times \widetilde{SU(2)}_R$. As before, since these

¹Relevance of Z_2 twist operators to the marginal deformations of the SCFT has earlier been discussed in [15, 16]

are top components of the short multiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ they can be added to the free SCFT as perturbations without violating the $\mathcal{N} = (4, 4)$ supersymmetry of the SCFT. The various quantum numbers of these operators are listed below.

Operator	$(j^3, \tilde{j}^3)_I$	$SU(2)_R \times \widetilde{SU(2)}_R$	$(h, \bar{h})$	
$\mathcal{T}_{(1)}^1 = T^{(\frac{1}{2}, \frac{1}{2})}$	$(0, 1)$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$	
$\mathcal{T}_{(0)}^1 = T^{(\frac{1}{2}, -\frac{1}{2})} + T^{(-\frac{1}{2}, \frac{1}{2})}$	$(0, 0)$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$	(2.45)
$\mathcal{T}_{(-1)}^1 = T^{(-\frac{1}{2}, -\frac{1}{2})}$	$(0, -1)$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$	
$\mathcal{T}^0 = T^{(-\frac{1}{2}, -\frac{1}{2})} - T^{(\frac{1}{2}, -\frac{1}{2})}$	$(0, 0)$	$(\mathbf{1}, \mathbf{1})$	$(1, 1)$	

The first three operators of the above table can be organized as a $(\mathbf{1}, \mathbf{3})$ under $SU(2)_I \times \widetilde{SU(2)}_I$. We will denote these 3 operators as $\mathcal{T}^1$. The last operator transforms as a scalar $(\mathbf{1}, \mathbf{1})$ under $SU(2)_I \times \widetilde{SU(2)}_I$ and is denoted by $\mathcal{T}^0$. The simplest way of figuring out the $(j^3, \tilde{j}^3)_I$ quantum numbers in the above table is to note that (a) the Σ -operators of (2.41) are singlets under $SU(2)_I \times \widetilde{SU(2)}_I$, as can be verified by computing the action on them of the operators I_1, I_2 and $\tilde{I}_1, \tilde{I}_2$, (b) the $\mathcal{T}$ -operators are obtained from Σ 's by the action of the supersymmetry currents as in (2.44) and (c) the quantum numbers of the supersymmetry currents under I_1, I_2 and $\tilde{I}_1, \tilde{I}_2$ are given by (2.25).

2.5.3 Higher twists

We now show that the twist operators corresponding to any other conjugacy class of $S(Q_1 Q_5)$ are irrelevant. Consider the class with $N_1 = Q_1 Q_5 - 3, N_3 = 1$ and the rest of $N_n = 0$. A representative element of this class is

$$(X_1 \rightarrow X_2, X_2 \rightarrow X_3, X_3 \rightarrow X_1), X_4 \rightarrow X_4, \dots, X_{Q_1 Q_5} \rightarrow X_{Q_1 Q_5}. \quad (2.46)$$

To make the action of this group element transparent we diagonalize the group action as follows.

$$\begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega^4 \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \\ X_3 \end{pmatrix} \quad (2.47)$$

where $\omega = \exp(2\pi i/3)$. These new coordinates are identified under the group action (2.46) $\phi_1 \rightarrow \phi_1, \phi_2 \rightarrow \omega^2 \phi_2$ and $\phi_3 \rightarrow \omega \phi_3$. These identifications are locally characteristic of the orbifold

$$R^4 \times R^4/\omega \times R^4/\omega^2 \quad (2.48)$$

The dimension of the supersymmetric twist operator which twists the coordinates by a phase $e^{2\pi i k/N}$ in 2 complex dimensions is $h(k, N) = k/N$ [12]. The twist operator which implements the action of the group element (2.46) combines the supersymmetric twist operators acting on ϕ_2 and ϕ_3 and therefore has total dimension

$$h = h(1, 3) + h(2, 3) = 1/3 + 2/3 = 1 \quad (2.49)$$

It is the superpartners of these which could be candidates for the blow up modes. However, these have weight $3/2$, These operators are therefore irrelevant.

For the class $N_1 = Q_1 Q_5 - k$, $N_k = k$ and the rest of $N_n = 0$, the total dimension of the twist operator is

$$h = \sum_{i=1}^{k-1} h(i, k) = (k-1)/2 \quad (2.50)$$

Its superpartner has dimension $k/2$. Now it is easy to see that all conjugacy classes other than the exchange of 2 elements give rise to irrelevant twist operators. Thus the orbifold $\mathcal{M}$ is resolved by the 4 blow up modes corresponding to the conjugacy class represented by (2.37). We have thus identified the 20 marginal operators of the $\mathcal{N} = (4, 4)$ SCFT on $\tilde{T}^4$. They are all top components of the $5(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ short multiplets. The $5(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ have 20 operators of conformal dimensions $(h, \bar{h}) = (1/2, 1/2)$. These are relevant operators for the SCFT. It would be interesting to investigate the role of these relevant operators.

2.6 The chiral primaries of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$

In this section we will explicitly construct all the chiral primaries corresponding to single particle states of the SCFT on the orbifold $\mathcal{M}$. For this purpose we will have to construct the twist operator corresponding to the conjugacy class $N_1 = Q_1 Q_5 - k$, $N_k = k$ and the rest of $N_n = 0$.

2.6.1 The k-cycle twist operator

We will extend the method of construction of the 2-cycle twist operator of Section 2.5.2 to the construction of the k-cycle twist operator. Consider the conjugacy class given by

$N_1 = Q_1 Q_5 - k$, $N_k = k$ and the rest of $N_n = 0$. A representative element of this class is the following group action

$$(X_1 \rightarrow X_2, \dots, X_k \rightarrow X_1), X_{k+1} \rightarrow X_{k+1}, \dots, X_{Q_1 Q_5} \rightarrow X_{Q_1 Q_5}. \quad (2.51)$$

We can diagonalize the group action as follows

$$\begin{pmatrix} \phi_k \\ \phi_{k-1} \\ \phi_{k-2} \\ \vdots \\ \phi_1 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{k-1} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(k-1)} \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 1 & \omega^{k-1} & \omega^{(k-1)2} & \dots & \omega^{(k-1)(k-1)} \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \\ X_3 \\ \vdots \\ X_k \end{pmatrix} \quad (2.52)$$

where $\omega = e^{2\pi i/k}$. These new coordinates are identified under the group action (2.51) as

$$\phi_1 \rightarrow \omega \phi_1, \quad \phi_2 \rightarrow \omega^2 \phi_2, \quad \phi_3 \rightarrow \omega^3 \phi_3, \quad \dots, \quad \phi_{k-1} \rightarrow \omega^{k-1} \phi_{k-1}, \quad \phi_k \rightarrow \omega^k \phi_k \quad (2.53)$$

These identifications are locally characteristic of the orbifold

$$R^4 \times R^4/\omega \times R^4/\omega^2 \times \dots \times R^4/\omega^{k-1} \quad (2.54)$$

The coordinate ϕ_m is twisted by the phase ω^m (m runs from $1 \dots k$). The bosonic twist operators corresponding to this twist are defined by the following OPE's

$$\begin{aligned} \partial \phi_m^1(z) \sigma_m^1(w, \bar{w}) &= \frac{\tau_m^1(w, \bar{w})}{(z-w)^{1-m/k}} & \partial \phi_m^{1\dagger}(z) \sigma_m^1(w, \bar{w}) &= \frac{\tau_m^{1\dagger}(w, \bar{w})}{(z-w)^{m/k}} \\ \partial \phi_m^2(z) \sigma_m^2(w, \bar{w}) &= \frac{\tau_m^2(w, \bar{w})}{(z-w)^{1-m/k}} & \partial \phi_m^{2\dagger}(z) \sigma_m^2(w, \bar{w}) &= \frac{\tau_m^{2\dagger}(w, \bar{w})}{(z-w)^{m/k}} \\ \bar{\partial} \tilde{\phi}_m^1(\bar{z}) \sigma_m^1(w, \bar{w}) &= \frac{\tilde{\tau}_m^1(w, \bar{w})}{(\bar{z}-\bar{w})^{m/k}} & \bar{\partial} \tilde{\phi}_m^{1\dagger}(\bar{z}) \sigma_m^1(w, \bar{w}) &= \frac{\tilde{\tau}_m^{1\dagger}(w, \bar{w})}{(\bar{z}-\bar{w})^{1-m/k}} \\ \bar{\partial} \tilde{\phi}_m^2(\bar{z}) \sigma_m^2(w, \bar{w}) &= \frac{\tilde{\tau}_m^2(w, \bar{w})}{(\bar{z}-\bar{w})^{m/k}} & \bar{\partial} \tilde{\phi}_m^{2\dagger}(\bar{z}) \sigma_m^2(w, \bar{w}) &= \frac{\tilde{\tau}_m^{2\dagger}(w, \bar{w})}{(\bar{z}-\bar{w})^{1-m/k}} \end{aligned} \quad (2.55)$$

As in Section 2.5.2 τ 's are excited twist operators. The fermionic twists are constructed from bosonized currents defined by

$$\begin{aligned} \chi_m^1(z) &= e^{iH_m^1(z)} & \chi_m^{1\dagger}(z) &= e^{-iH_m^1(z)} \\ \chi_m^2(z) &= e^{iH_m^2(z)} & \chi_m^{2\dagger}(z) &= e^{-iH_m^2(z)} \end{aligned} \quad (2.56)$$

Where the χ_m 's are the superpartners of the bosons ϕ_m 's. The twist operators corresponding to the fermions χ_m 's are given by $e^{\pm imH_m/k}$.

We now assemble all these operators to construct the k-cycle twist operator which is a chiral primary. The k-cycle twist operator is given by

$$\Sigma_{(12\dots k)}^{(k-1)/2} = \prod_{m=1}^{k-1} \left[\sigma_m^1(z, \bar{z}) \sigma_m^2(z, \bar{z}) e^{imH_m^1(z)/k} e^{-imH_m^2(z)/k} e^{im\tilde{H}_m^1(\bar{z})/k} e^{-im\tilde{H}_m^2(\bar{z})/k} \right] \quad (2.57)$$

The subscript $(12\dots k)$ refers to the fact that these twist operators were constructed for the representative group element (2.51) which cyclically permutes the $1, \dots, k$ labels of the coordinates of $\tilde{T}^4$. The superscript $(k-1)/2$ stands for the conformal dimension of this operator. As we saw in Section 2.5.3 the conformal dimension of the twist operator for the conjugacy class $N_1 = Q_1Q_5 - k$, $N_k = k$ and the rest of $N_n = 0$ is $(h, \bar{h}) = ((k-1)/2, (k-1)/2)$. The twist operator for the conjugacy class under consideration is obtained by summing over the k-cycle twist operators for all representative element of these class.

$$\Sigma^{(k-1)/2}(z, \bar{z}) = \sum_{\{i_1, \dots, i_k\}} \Sigma_{i_1 i_2 \dots i_k}(z, \bar{z}) \quad (2.58)$$

where the sum runs over all k -tuples $\{i_1, \dots, i_k\}$ such that $i_1 \neq i_2 \neq \dots \neq i_k$. i_m take values from 1 to Q_1Q_5 . The operator $\Sigma^{(k-1)/2}$ is a chiral primary with conformal dimension $(h, \bar{h}) = ((k-1)/2, (k-1)/2)$ and $(j_R^3, \tilde{j}_R^3) = ((k-1)/2, (k-1)/2)$. As the largest cycle is of length Q_1Q_5 , the maximal dimension of the k-cycle twist operator is $((Q_1Q_5 - 1)/2, (Q_1Q_5 - 1)/2)$. It belongs to the bottom component of the short multiplet $(\mathbf{k}, \mathbf{k})_{\mathbf{S}}$. The other components of the shortmultiplet $(\mathbf{k}, \mathbf{k})_{\mathbf{S}}$ corresponding to the k-cycle twists can be generated by the action of supersymmetry currents and the R-symmetry currents of the $\mathcal{N} = (4, 4)$ theory on $\mathcal{M}$.

2.6.2 The complete set of chiral primaries

We have seen in Section 2.5 there are five chiral primaries corresponding to the shortmultiplet $5(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$. In this section we will construct the complete set of chiral primaries from single particle states of the SCFT on $\mathcal{M}$. It is known that the chiral primaries with weight $(h, \bar{h})$ of a $\mathcal{N} = (4, 4)$ superconformal field theory on a manifold K correspond to the elements of the cohomology $\mathcal{H}_{2h, 2\bar{h}}(K)$ [13]. The chiral primaries are formed by the product of the chiral primaries corresponding to the cohomology of the diagonal $\tilde{T}^4$ denoted by B^4 (the sum of

all copies of $\tilde{T}^4$) and the various k-cycle chiral primaries constructed in Section 2.6.1. We will list the chiral primaries below

Chiral primaries with $h - \bar{h} = 0$

All the k-cycle chiral primaries have $h - \bar{h} = 0$. To construct chiral primaries with $h - \bar{h} = 0$ we need the four chiral primaries which correspond to the cohomology $\mathcal{H}_{11}(B^4)$ with weight $(1/2, 1/2)$. They are given in (2.33). Using this we can construct the following chiral primaries

$$\begin{aligned} \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^1(z) \tilde{\Psi}_A^1(\bar{z}) & \quad \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^1(z) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \\ \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) & \quad \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \end{aligned} \quad (2.59)$$

where summation over A is implied. These four operators have conformal dimension $(k/2, k/2)$. There is one more chiral primary corresponding to the cohomology $\mathcal{H}_{22}(B^4)$ for which $h - \bar{h} = 0$. It is given by

$$\Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \quad (2.60)$$

where summation over all indices of A is implied. This chiral primary corresponds to the top form of B^4 . The cohomology $\mathcal{H}_{00}(B^4)$ gives rise to a chiral primaries of conformal dimension $(k/2, k/2)$. It is given by

$$\Sigma^{(k-2)/2}(z, \bar{z}) \Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \quad (2.61)$$

From the equation above we see that these chiral primaries exist only of $k \geq 2$. Finally we have the chiral primary $\Sigma^{(k)/2}(z, \bar{z})$ of conformal dimension $(k/2, k/2)$. Thus for $k \geq 2$ and $k \leq Q_1 Q_5 - 1$ there are 6 chiral primaries of dimension $(k/2, k/2)$

The complete list of chiral primaries with $(h, \bar{h})$ with $h - \bar{h} = 0$ corresponding to single particle states are given by

$(h, \bar{h})$	Degeneracy	
$(1/2, 1/2)$	5	
$(1, 1)$	6	
$(3/2, 3/2)$	6	
$\vdots$	$\vdots$	
$((Q_1 Q_5 - 1)/2, (Q_1 Q_5 - 1)/2)$	6	
$((Q_1 Q_5)/2, (Q_1 Q_1)/2)$	5	
$((Q_1 Q_5 + 1)/2, (Q_1 Q_5 + 1)/2)$	1	

(2.62)

In the above table we have ignored the vacuum with weight $(h, \bar{h}) = (0, 0)$.

Chiral primaries with $h - \bar{h} = 1/2$

The chiral primaries of B^4 which correspond to the elements of the cohomology $\mathcal{H}_{10}(B^4)$ are given by

$$\sum_{A=1}^{Q_1 Q_5} \Psi_A^1(z) \quad \text{and} \quad \sum_{A=1}^{Q_1 Q_5} \Psi_A^{2\dagger}(z) \quad (2.63)$$

We can construct chiral primaries with weight $((k+1)/2, k/2)$ by taking the product of the above chiral primaries with the twist operator $\Sigma^{k/2}(z, \bar{z})$. These give the following chiral primaries

$$\Sigma^{k/2}(z, \bar{z}) \sum_{A=1}^{Q_1 Q_5} \Psi_A^1(z) \quad \text{and} \quad \Sigma^{k/2}(z, \bar{z}) \sum_{A=1}^{Q_1 Q_5} \Psi_A^{2\dagger}(z) \quad (2.64)$$

The chiral primary of the diagonal B^4 which correspond to the elements of the cohomology $\mathcal{H}_{21}(B^4)$ are

$$\Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) \quad \text{and} \quad \Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \quad (2.65)$$

Here summation over all the three indices of A is implied. From these the one can construct chiral primaries with weight $((k+1)/2, k/2)$ are follows

$$\Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) \quad \text{and} \quad \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^1(z) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \quad (2.66)$$

Therefore there are 4 chiral primaries with weight $((k+1)/2, k/2)$ for $1 \leq k \leq (Q_1 Q_5 - 1)$ and 2 chiral primaries with weight $((Q_1 Q_5 + 1)/2, Q_1 Q_5/2)$. There are also 2 chiral primaries with weight $(1/2, 0)$.

Chiral primaries with $\bar{h} - h = 1/2$

The procedure for constructing these chiral primaries are identical to the case $h - \bar{h} = 1/2$. The four chiral primaries with weight $(k/2, (k+1)/2)$ are given by

$$\begin{aligned} \Sigma^{k/2}(z, \bar{z}) \sum_{A=1}^{Q_1 Q_5} \tilde{\Psi}_A^1(\bar{z}) & \quad \Sigma^{k/2}(z, \bar{z}) \sum_{A=1}^{Q_1 Q_5} \tilde{\Psi}_A^{2\dagger}(\bar{z}) \\ \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^1(z) \tilde{\Psi}_A^1(\bar{z}) \tilde{\Psi}_A^{2\dagger}(\bar{z}) & \quad \Sigma^{(k-1)/2}(z, \bar{z}) \Psi_A^{2\dagger}(z) \tilde{\Psi}_A^1(\bar{z}) \tilde{\Psi}_A^{2\dagger}(\bar{z}) \end{aligned} \quad (2.67)$$

There are 4 chiral primaries with weight $(k/2, (k+1)/2)$ for $1 \leq k \leq (Q_1 Q_5 - 1)$ 2 chiral primaries with weight $(Q_1 Q_5/2, (Q_1 Q_5 + 1)/2)$ and 2 chiral primaries with weight $(0, 1/2)$.

Chiral primaries with $h - \bar{h} = 1$

As in the previous cases let us first look at the chiral primaries corresponding to the cohomology element $\mathcal{H}_{20}(B^4)$. There is only one element which is given by

$$\Psi_A^1(z)\Psi_A^{2\dagger}(z) \quad (2.68)$$

where summation over A is implied. There is a single chiral primary with weight $((k + 2)/2, k/2)$ constructed out of the above chiral primary is

$$\Sigma^{k/2}(z, \bar{z})\Psi_A^1(z)\Psi_A^{2\dagger}(z) \quad (2.69)$$

Thus there is a one chiral primary with weight $((k + 2)/2, k/2)$ for $0 \leq k \leq (Q_1 Q_5 - 1)$

The operator product expansion of two chiral primaries will give rise to other chiral primaries consistent with conservation laws. There are known to form a ring. It will be interesting to understand the structure of this ring.

Chiral primaries with $\bar{h} - h = 1$

The construction of these is parallel to the case for $h - \bar{h} = 1$. The single chiral primary with weight $(k/2, (k + 2)/2)$ for $0 \leq k \leq (Q_1 Q_5 - 1)$ is given by

$$\Sigma^{k/2}(z, \bar{z})\tilde{\Psi}_A^1(\bar{z})\tilde{\Psi}_A^{2\dagger}(\bar{z}) \quad (2.70)$$

2.7 Shortmultiplets of $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$

Using the results of Section 2.6 we will write the complete set of shortmultiplets of single particle states of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$. In Chapter 3 we will compare this set of shortmultiplets with that obtained from supergravity. We will see in Chapter 3 that supergravity is a good approximation in string theory only when $Q_1 \rightarrow \infty, Q_5 \rightarrow \infty$. Therefore we write down the list of shortmultiplets for $(\tilde{T}^4)^{(\infty)}/S(\infty)$. Basically this means that the list of chiral primaries of the previous Section 2.6 does not terminate.

We have seen that each chiral primary of weight (h, h') gives rise to the short multiplet $(2\mathbf{h} + \mathbf{1}, 2\mathbf{h}' + \mathbf{1})_S$. Therefore the results of Section 2.6 indicate that the list of shortmultiplets corresponding to the single particle states of $\mathcal{N} = (4, 4)$ SCFT on $(\tilde{T}^4)^{(\infty)}/S(\infty)$.

is given by

$$\begin{aligned}
& 5(\mathbf{2}, \mathbf{2})_S + 6 \oplus_{\mathbf{m} \geq \mathbf{3}} (\mathbf{m}, \mathbf{m})_S \\
& 2(\mathbf{1}, \mathbf{2})_S + 2(\mathbf{2}, \mathbf{1})_S + (\mathbf{1}, \mathbf{3})_S + (\mathbf{3}, \mathbf{1})_S \\
& \oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m}, \mathbf{m} + \mathbf{2})_S + (\mathbf{m} + \mathbf{2}, \mathbf{m})_S + 4(\mathbf{m}, \mathbf{m} + \mathbf{1})_S + 4(\mathbf{m} + \mathbf{1}, \mathbf{m})_S]
\end{aligned} \tag{2.71}$$

2.8 The location of the symmetric product

The complete specification of the D1/D5 system includes various moduli. Most of the studies of the D1/D5 system so far have been focused on the situation with no moduli. It is known from supergravity that the D1/D5 system with no moduli turned on is marginally stable with respect to the decay of a subsystem consisting of Q'_5 D5 and Q'_1 D1-branes. It has been observed recently [17] that such a decay in fact signals a singularity in the world volume gauge theory associated with the origin of the Higgs branch. The issue of stability in supergravity in the context of the D1/D5 system has also been discussed in [6, 18].

The singularity mentioned above leads to a singular conformal field theory and hence to a breakdown of string perturbation theory. However, generic values of the supergravity moduli which do not involve fragmentation into constituents are described by well-defined conformal field theories and therefore string perturbation theory makes sense. We have seen in section 2.5 that important singularity structure of the $\mathcal{M} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$ is locally of the type R^4/Z_2 . The resolution of this singularity gives rise to marginal operators. An orbifold theory realized as a free field SCFT on R^4/Z_2 is nonsingular as all correlations functions are finite. The reason for this can be understood from the linear sigma model description of the R^4/Z_2 singularity discussed in section 2.9.1. We will see that though the R^4/Z_2 singularity is geometrically singular the SCFT is finite because it corresponds to a non-zero theta term in the linear sigma model. The geometric resolution of this singularity corresponds to adding Fayet-Iliopoulos terms to the D-term equations of the linear sigma model. This deforms the R^4/Z_2 singularity to an Eguchi-Hansen space. In the orbifold theory this deformation is caused by the twist operator $\mathcal{T}^1$. The Eguchi-Hansen space is asymptotically R^4/Z_2 but the singularity at the origin is blown up to a 2-sphere. One can use the $SU(2)_R$ symmetry of the linear sigma model to rotate the three Fayet-Iliopoulos

terms to one term. This term corresponds to the radius of the blown up 2-sphere. The theta term of the linear sigma model corresponds to B -flux through the 2-sphere. The change of this B -flux is caused by deforming the orbifold SCFT by the twist operator $\mathcal{T}^0$. Thus SCFT realized as a free field theory on the orbifold R^4/Z_2 is regular even though the 2-sphere is squashed to zero size because of the non-zero value of B -flux trapped in the squashed 2-sphere [19].

The realization of the SCFT of the D1/D5 system as $\mathcal{N} = (4, 4)$ theory on $\mathcal{M}$ implies that we are at a point in the moduli space of the D1/D5 system at which the orbifold is geometrically singular but because of the non-zero value of the theta term the SCFT is regular and *not at the singularity corresponding to fragmentation*. In other words, the orbifold SCFT corresponds to a bound state of Q_1 D1-branes and Q_5 D5-branes (henceforth denoted as the (Q_1, Q_5) bound state). Thus we use the free field realization of $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$ and its resolutions using the marginal operators of this theory to describe the boundary SCFT corresponding to the D1/D5 system at generic values of the moduli.

The symmetric product orbifold $\mathcal{M}$ as we have seen before in section 2.2 is the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on T^4 . We argue that the orbifold $\mathcal{M}$ resolved by the twist operators $\mathcal{T}^1$ is the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on a non-commutative torus T^4 . We first look at the case of instantons on R^4 . The moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory is given by the D-terms of a 1 + 1 dimensional $U(Q_1)$ gauge theory with $(4, 4)$ supersymmetry and Q_5 hypermultiplets. On deforming these D-term equations by Fayet-Iliopoulos terms we obtain the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on a non-commutative R^4 [20]. Now let us turn to instantons on T^4 . The R^4/Z_2 singularity of the orbifold $\mathcal{M}$ is resolved by adding Fayet-Iliopoulos terms to the D-term equations in the linear sigma model description of the R^4/Z_2 singularity. As we have seen in the orbifold theory this is caused by adding the twist operators $\mathcal{T}^1$ to the free theory. The analogy with the case of instantons on R^4 gives us the hint that the resolved orbifold $\mathcal{M}$ is the moduli space of Q_1 instantons of a $U(Q_5)$ gauge theory on a non-commutative torus T^4 . It will be interesting to make this argument more precise.

2.9 The linear sigma model

In this section we will analyze the gauge theory description of the D1/D5 system. We show that the gauge theory has four parameters which control the break up of the (Q_1, Q_5) system to subsystems (Q'_1, Q'_5) and (Q''_1, Q''_5) . These are the Fayet-Iliopoulos D-terms and the theta term in the effective $U(1)$ $(4, 4)$ linear sigma model of the relative coordinate between the subsystems (Q'_1, Q'_5) and (Q''_1, Q''_5) . To motivate this we will review the linear sigma model corresponding to the of the R^4/Z_2 singularity.

2.9.1 The linear sigma model description of R^4/Z_2

The linear sigma model is a $1 + 1$ dimensional $U(1)$ gauge theory with $(4, 4)$ supersymmetry [21]. It has 2 hypermultiplets charged under the $U(1)$. The scalar fields of the hypermultiplets can be organized as doublets under the $SU(2)_R$ symmetry of the $(4, 4)$ theory as

$$\chi_1 = \begin{pmatrix} A_1 \\ B_1^\dagger \end{pmatrix} \text{ and } \chi_2 = \begin{pmatrix} A_2 \\ B_2^\dagger \end{pmatrix} \quad (2.72)$$

The A 's have charge $+1$ and the B 's have charge -1 under the $U(1)$. The vector multiplet has 4 real scalars φ_i , $i = 1, \dots, 4$. They do not transform under the $SU(2)_R$. One can include 4 parameters in this theory consistent with $(4, 4)$ supersymmetry. They are the 3 Fayet-Iliopoulos terms and the theta term.

Let us first investigate the hypermultiplet moduli space of this theory with the 3 Fayet-Iliopoulos terms and the theta term set to zero. The Higgs phase of this theory is obtained by setting ϕ_i and the D-terms to zero. The D-term equations are

$$|A_1|^2 + |A_2|^2 - |B_1|^2 - |B_2|^2 = 0 \quad (2.73)$$

$$A_1 B_1 + A_2 B_2 = 0$$

The hypermultiplet moduli space is the space of solutions of the above equations modded out by the $U(1)$ gauge symmetry. Counting the number of degrees of freedom indicate that this space is 4 dimensional. To obtain the explicit form of this space it is convenient to introduce the following gauge invariant variables

$$M = A_1 B_2 \quad N = A_2 B_1 \quad (2.74)$$

$$P = A_1 B_1 = -A_2 B_2 \quad (2.75)$$

These variables are not independent. Setting the D-terms equal to zero and modding out the resulting space by $U(1)$ is equivalent to the equation

$$P^2 + MN = 0. \quad (2.76)$$

This homogeneous equation is an equation of the space R^4/Z_2 . To see this the solution of the above equation can be parameterized by 2 complex numbers (ζ, η) such that

$$P = i\zeta\eta \quad M = \zeta^2 \quad N = \eta^2 \quad (2.77)$$

Thus the point (ζ, η) and $(-\zeta, -\eta)$ are the same point in the space of solutions of (2.76). We have shown that the hypermultiplet moduli space is R^4/Z_2 .

The above singularity at the origin of the moduli space is a geometric singularity in the hypermultiplet moduli space. We now argue that this singularity is a genuine singularity of the SCFT that the linear sigma model flows to in the infrared. At the origin of the classical moduli space the Coulomb branch meets the Higgs branch. In addition to the potential due to the D-terms the linear sigma model contains the following term in the superpotential²

$$V = (|A_1|^2 + |A_2|^2 + |B_1|^2 + |B_2|^2)(\varphi_1^2 + \varphi_2^2 + \varphi_3^2 + \varphi_4^2) \quad (2.78)$$

Thus at the origin of the hypermultiplet moduli space a flat direction for the Coulomb branch opens up. The ground state at this point is not normalizable due to the non-compactness of the Coulomb branch. This renders the infrared SCFT singular.

This singularity can be avoided in two distinct ways. If one turns on the Fayet-Iliopoulos D-terms, the D-term equations are modified to [21]

$$|A_1|^2 + |A_2|^2 - |B_1|^2 - |B_2|^2 = r_3 \quad (2.79)$$

$$A_1 B_1 + A_2 B_2 = r_1 + ir_2$$

²These terms can be understood from the coupling $A_\mu A^\mu \chi^* \chi$ in six dimensions, and recognizing that under dimensional reduction to two dimensions φ_i 's appear from the components of A_μ in the compact directions.

Where r_1, r_2, r_3 are the 3 Fayet-Iliopoulos D-terms transforming as the adjoint of the $SU(2)_R$. Now the origin is no more a solution of these equations and the non-compactness of the Coulomb branch is avoided. In this case wave-functions will have compact support all over the hypermultiplet moduli space. This ensures that the infrared SCFT is non-singular. Turning on the Fayet-Iliopoulos D-terms thus correspond to the geometric resolution of the singularity. The resolved space is known to be [21, 19] described by an Eguchi-Hanson metric in which $r_{1,2,3}$ parameterize a shrinking two-cycle.

The second way to avoid the singularity in the SCFT is to turn on the theta angle θ . This induces a constant electric field in the vacuum. This electric field is screened at any other point than the origin in the hypermultiplet moduli space as the $U(1)$ gauge field is massive with a mass proportional to the vacuum expectation value of the hypers. At the origin the $U(1)$ field is not screened and thus it contributes to the energy density of the vacuum. This energy is proportional to θ^2 . Thus turning on the theta term lifts the flat directions of the Coulomb branch. This ensures that the corresponding infrared SCFT is well defined though the hypermultiplet moduli space remains geometrically singular. In terms of the Eguchi-Hanson space, the θ -term corresponds to a flux of the antisymmetric tensor through the two-cycle mentioned above.

The $(4, 4)$ SCFT on R^4/Z_2 at the orbifold point is well defined. Since the orbifold has a geometric singularity but the SCFT is non-singular it must correspond to the linear sigma model with a finite value of θ and the Fayet-Iliopoulos D-terms set to zero. Deformations of the R^4/Z_2 orbifold by its 4 blow up modes correspond to changes in the Fayet-Iliopoulos D-terms and theta term of the linear sigma model³ The global description of the moduli of a $\mathcal{N} = (4, 4)$ SCFT on a resolved R^4/Z_2 orbifold is provided by the linear sigma model. In conclusion let us describe this linear sigma model in terms of the gauge theory of D-branes. The theory described above arises on a single D1-brane in presence of 2 D5-branes. The singularity at the point $r_1, r_2, r_3, \theta = 0$ is due to noncompactness of the flat direction of the Coulomb branch. Thus it corresponds to the physical situation of the D1-brane leaving the

³If we identify the $SU(2)_R$ of the linear sigma-model with $\widetilde{SU(2)}_I$ of the orbifold SCFT, then the Fayet-Iliopoulos parameters will correspond to $\mathcal{T}^1$ and the θ -term to $\mathcal{T}^0$. This is consistent with Witten's observation [22] that $SO(4)_E$ symmetry of the linear sigma-model (one that rotates the ϕ_i 's) corresponds to the $SU(2)_R$ of the orbifold SCFT.

D5-branes.

2.9.2 The gauge theory description of the moduli of the D1/D5 system

As we have seen in Section 2 the resolutions of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ is described by 4 marginal operators which were identified in the last subsection with the Fayet-Iliopoulos D-terms and the theta term of the linear sigma model description of the R^4/Z_2 singularity. We want to now indicate how these four parameters would make their appearance in the gauge theory description of the full D1/D5 system.

Motivated by the D-brane description of the R^4/Z_2 singularity we look for the degrees of freedom characterizing the break up of (Q_1, Q_5) system to (Q'_1, Q'_5) and (Q''_1, Q''_5) . Physically the relevant degree of freedom describing this process is the relative coordinate between the centre of mass of the (Q'_1, Q'_5) system and the (Q''_1, Q''_5) . We will describe the effective theory of this degree of freedom below.

For the bound state (Q_1, Q_5) the hypermultiplets the $\chi_{a,\bar{b}}$ are charged under the relative $U(1)$ of $U(Q_1) \times U(Q_5)$, that is under the gauge field $A_\mu = Tr_{U(Q_1)}(A_\mu^{a\bar{a}}) - Tr_{U(Q_5)}(A_\mu^{b\bar{b}})$. The gauge multiplet corresponding to the relative $U(1)$ corresponds to the degree of freedom of the relative coordinate between the centre of mass of the collection of Q_1 D1-branes and Q_5 D5-branes. At a generic point of the Higgs phase, all the $\chi_{a\bar{b}}$'s have expectation values, thus making this degree of freedom becomes massive. This is consistent with the fact that we are looking at the bound state (Q_1, Q_5) .

Consider the break up of the (Q_1, Q_5) bound state to the bound states (Q'_1, Q'_5) and (Q''_1, Q''_5) . To find out the charges of the hypermultiplets under the various $U(1)$, we will organize the hypers as

$$\begin{pmatrix} \chi_{a'b'} & \chi_{a'b''} \\ \chi_{a''b'} & \chi_{a''b''} \end{pmatrix}, \quad \begin{pmatrix} Y_{i(a'\bar{a}')}^{(1)} & Y_{i(a'\bar{a}'')}^{(1)} \\ Y_{i(a''\bar{a}')}^{(1)} & Y_{i(a''\bar{a}'')}^{(1)} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} Y_{i(b'\bar{b}')}^{(5)} & Y_{i(b'\bar{b}'')}^{(5)} \\ Y_{i(b''\bar{b}')}^{(5)} & Y_{i(b''\bar{b}'')}^{(5)} \end{pmatrix} \quad (2.80)$$

where $a', \bar{a}'$ runs from $1, \dots, Q'_1$, $b', \bar{b}'$ from $1, \dots, Q'_5$, $a'' \bar{a}''$ from $1, \dots, Q''_1$ and $b'', \bar{b}''$ from $1, \dots, Q''_5$. We organize the scalars of the vector multiplet corresponding to the gauge group

$U(Q_1)$ and $U(Q_5)$ as

$$\begin{pmatrix} \phi_i^{(1)a'\bar{a}'} & \phi_i^{(1)a'\bar{a}''} \\ \phi_i^{(1)a''\bar{a}'} & \phi_i^{(2)a'\bar{a}''} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \phi_i^{(5)b'\bar{b}'} & \phi_i^{(5)b'\bar{b}''} \\ \phi_i^{(5)b''\bar{b}'} & \phi_i^{(5)b'\bar{b}''} \end{pmatrix} \quad (2.81)$$

where $i = 1, 2, 3, 4$.

Let us call the the $U(1)$ gauge fields (traces) of $U(Q'_1)$, $U(Q'_5)$, $U(Q''_1)$, $U(Q''_5)$ as A'_1 , A'_5 , A''_1 , A''_5 respectively. We will also use the notation $A'_\pm \equiv A'_1 \pm A'_5$ and $A''_\pm \equiv A''_1 \pm A''_5$.

As we are interested in the bound states (Q'_1, Q'_5) and (Q''_1, Q''_5) , in what follows we will work with a specific classical background in which we give vev 's to the block-diagonal hypers $\chi_{a'b'}$, $\chi_{a''b''}$, $Y_{i(a'\bar{a}')}^{(1)}$, $Y_{i(b'\bar{b}')}^{(5)}$, $Y_{i(a''\bar{a}'')}^{(1)}$ and $Y_{i(b''\bar{b}'')}^{(5)}$. These vev 's are chosen so that the classical background satisfies the D-term equations [3].

The vev 's of the χ 's render the fields A'_- and A''_- massive with a mass proportional to vev 's. In the low energy effective Lagrangian these gauge fields can therefore be neglected. In the following we will focus on the $U(1)$ gauge field $A_r = 1/2(A'_+ - A''_+)$ which does not get mass from the above vev 's. The gauge multiplet corresponding to A_r contains four real scalars denoted below by φ_i . These represent the relative coordinate between the centre of mass of the (Q'_1, Q'_5) and the (Q''_1, Q''_5) bound states. We will be interested in the question whether the φ_i 's remain massless or otherwise. The massless case would correspond to a non-compact Coulomb branch and eventual singularity of the SCFT.

In order to address the above question we need to find the low energy degrees of freedom which couple to the gauge multiplet corresponding to A_r .

The fields charged under A_r are the hypermultiplets $\chi_{a'\bar{b}''}$, $\chi_{a''\bar{b}'}$, $Y_{i(a'\bar{a}'')}^{(1)}$, $Y_{i(a''\bar{a}')}^{(1)}$, $Y_{i(b'\bar{b}'')}^{(5)}$, $Y_{i(b''\bar{b}')}^{(5)}$ and the vector multiplets $\phi_i^{(1)a'\bar{a}''}$, $\phi_i^{(1)a''\bar{a}'}$, $\phi_i^{(5)b'\bar{b}''}$, $\phi_i^{(5)b''\bar{b}'}$. In order to find out which of these are massless, we look at the following terms in the Lagrangian of $U(Q_1) \times U(Q_5)$ gauge theory:

$$L = L_1 + L_2 + L_3 + L_4 \quad (2.82)$$

$$L_1 = \chi_{a_1\bar{b}_1}^* \phi_i^{(1)a_2\bar{a}_1*} \phi_i^{(1)a_2\bar{a}_3} \chi_{a_3\bar{b}_1}$$

$$L_2 = \chi_{a_1\bar{b}_1}^* \phi_i^{(5)b_1\bar{b}_2*} \phi_i^{(5)b_3\bar{b}_2} \chi_{a_1\bar{b}_3}$$

$$L_3 = Tr([Y_i^{(1)}, Y_j^{(1)}][Y_i^{(1)}, Y_j^{(1)}])$$

$$L_4 = Tr([Y_i^{(5)}, Y_j^{(5)}][Y_i^{(5)}, Y_j^{(5)}])$$

where the a_i 's run from $1, \dots, Q_1$ and the b_i 's run from $1, \dots, Q_5$. The terms L_1 and L_2 originate from terms of the type $|A_M \chi|^2$ where $A_M \equiv (A_\mu, \phi_i)$ is the $(4, 4)$ vector multiplet in two dimensions. The terms L_3 and L_4 arise from commutators of gauge fields in compactified directions.

The fields Y are in general massive. The reason is that the traces $y_i'^{(1)} \equiv Y_{i(a'\bar{a}')}^{(1)}$, representing the centre-of-mass position in the T^4 of Q'_1 D1-branes, and $y_i''^{(1)} \equiv Y_{i(a''\bar{a}'')}^{(1)}$, representing the centre-of-mass position in the T^4 of Q''_1 D1-branes, are neutral and will have *vev*'s which are generically separated (the centres of mass can be separated in the torus even when they are on top of each other in physical space). The mass of $Y_{i(a'\bar{a}'')}^{(1)}, Y_{i(a''\bar{a}')}^{(1)}$ can be read off from the term L_3 in (2.82), to be proportional to $(y'^{(1)} - y''^{(1)})^2$. Similarly the mass of $Y_{i(b'\bar{b}'')}^{(5)}, Y_{i(b''\bar{b}')}^{(5)}$ is proportional to $(y'^{(5)} - y''^{(5)})^2$ (as can be read off from the term L_4 in (2.82)) where $y'^{(5)}$ and $y''^{(5)}$ are the centers of mass of the Q'_5 D5-branes and Q''_5 D5-branes along the direction of the dual four torus $\hat{T}^4$. (At special points when their centres of mass coincide, these fields become massless. The analysis for these cases can also be carried out by incorporating these fields in (2.85)-(2.87), with no change in the conclusion) The fields $\phi_i^{(1)a'\bar{a}'}, \phi_i^{(1)a''\bar{a}'}$ are also massive. Their masses can be read off from the L_1 in (2.82). Specifically they arise from the following terms

$$\chi_{a''\bar{b}'}^* \phi_i^{(1)a'\bar{a}''*} \phi_i^{(1)a'\bar{a}''} \chi_{a''\bar{b}''} + \chi_{a'\bar{b}'}^* \phi_i^{(1)a''\bar{a}''*} \phi_i^{(1)a''\bar{a}''} \chi_{a'\bar{b}''} \quad (2.83)$$

where a'_i run from $1, \dots, Q'_1$ and a''_i run from $1, \dots, Q''_1$. These terms show that their masses are proportional to the expectation values of the hypers $\chi_{a'b'}$ and $\chi_{a''b''}$. Similarly the terms of L_2 in (2.82)

$$\chi_{a''\bar{b}_1'}^* \phi_i^{(5)b'_1\bar{b}''*} \phi_i^{(5)b'_1\bar{b}''} \chi_{a''\bar{b}_2''} + \chi_{a'\bar{b}_1'}^* \phi_i^{(5)b'_1\bar{b}''*} \phi_i^{(5)b'_1\bar{b}''} \chi_{a'\bar{b}_2''} \quad (2.84)$$

show that the fields $\phi_i^{(5)b'\bar{b}''} \phi_i^{(5)b''\bar{b}'}$ are massive with masses proportional to the expectation values of the hypers $\chi_{a'\bar{a}''}$ and $\chi_{a''\bar{b}''}$. In the above equation b'_i take values from $1, \dots, Q'_5$ and b''_i take values from $1, \dots, Q''_5$. Note that these masses remain non-zero even in the limit when the (Q'_1, Q'_5) and (Q''_1, Q''_5) are on the verge of separating.

Thus the relevant degrees of freedom describing the splitting process is a $1+1$ dimensional $U(1)$ gauge theory of A_r with $(4, 4)$ supersymmetry. The matter content of this theory

consists of hypermultiplets $\chi_{a'\bar{b}''}$ with charge +1 and $\chi_{a''\bar{b}'}$ with charge -1.

Let us now describe the dynamics of the splitting process. This is given by analyzing the hypermultiplet moduli space of the effective theory described above with the help of the D-term equations:

$$\begin{aligned} A_{a'\bar{b}''}A_{a''\bar{b}'}^* - A_{a''\bar{b}'}A_{a'\bar{b}''}^* - B_{b''\bar{a}'}B_{b'\bar{a}''}^* + B_{b'\bar{a}''}B_{b''\bar{a}'}^* &= 0 \\ A_{a'\bar{b}''}B_{b''\bar{a}'} - A_{a''\bar{b}'}B_{b'\bar{a}''} &= 0 \end{aligned} \quad (2.85)$$

In the above equations the sum over a', b', a'', b'' is understood. These equations are generalized version of (2.73) discussed for the R^4/Z_2 singularity in Section 4.1. At the origin of the Higgs branch where the classical moduli space meets the Coulomb branch this linear sigma model would flow to an infrared conformal field theory which is singular. The reason for this is the same as for the R^4/Z_2 case. The linear sigma model contains the following term in the superpotential

$$V = (A_{a'\bar{b}''}A_{a''\bar{b}'}^* + A_{a''\bar{b}'}A_{a'\bar{b}''}^* + B_{b''\bar{a}'}B_{b'\bar{a}''}^* + B_{b'\bar{a}''}B_{b''\bar{a}'}^*)(\varphi_1^2 + \varphi_2^2 + \varphi_3^2 + \varphi_4^2) \quad (2.86)$$

As in the discussion of the R^4/Z_2 case, at the origin of the hypermultiplet moduli space the flat direction of the Coulomb branch leads to a ground state which is not normalizable. This singularity can be avoided by deforming the D-term equations by the Fayet-Iliopoulos terms:

$$\begin{aligned} A_{a'\bar{b}''}A_{a''\bar{b}'}^* - A_{a''\bar{b}'}A_{a'\bar{b}''}^* - B_{b''\bar{a}'}B_{b'\bar{a}''}^* + B_{b'\bar{a}''}B_{b''\bar{a}'}^* &= r_3 \\ A_{a'\bar{b}''}B_{b''\bar{a}'} - A_{a''\bar{b}'}B_{b'\bar{a}''} &= r_1 + ir_3 \end{aligned} \quad (2.87)$$

We note here that the Fayet-Iliopoulos terms break the relative $U(1)$ under discussion and the gauge field becomes massive. The reason is that the D-terms with the Fayet-Iliopoulos do not permit all A, B 's in the above equation to simultaneously vanish. At least one of them must be non-zero. As these A, B 's are charged under the $U(1)$, the non-zero value of A, B gives mass to the vector multiplet. This can be seen from the potential (2.86). The scalars of the vector multiplet become massive with the mass proportional to the vev 's of A, B . Thus the relative $U(1)$ is broken.

The singularity associated with the non-compact Coulomb branch can also be avoided by turning on the θ term, the mechanism being similar to the one discussed in the previous

subsection. If any of the 3 Fayet-Iliopoulos D-terms or the θ term is turned on, the flat directions of the Coulomb branch are lifted, leading to normalizable ground state is of the Higgs branch. This prevents the breaking up of the (Q_1, Q_5) system to subsystems. Thus we see that the 4 parameters which resolve the singularity of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ make their appearance in the gauge theory as the Fayet-Iliopoulos terms and the theta term.

It would be interesting to extract the singularity structure of the the gauge theory of the D1/D5 system through mappings similar to (2.74)- (2.77)⁴.

2.9.3 The case $(Q_1, Q_5) \rightarrow (Q_1 - 1, Q_5) + (1, 0)$: splitting of 1 D1-brane

It is illuminating to consider the special case in which 1 D1-brane splits off from the bound state (Q_1, Q_5) . The effective dynamics is again described in terms of a $U(1)$ gauge theory associated with the relative separation between the single D1-brane and the bound state $(Q_1 - 1, Q_5)$. The massless hypermultiplets charged under this $U(1)$ correspond to open strings joining the single D1-brane with the D5-branes and are denoted by

$$\chi_{b'} = \begin{pmatrix} A_{b'} \\ B_{b'}^\dagger \end{pmatrix} \quad (2.88)$$

The D-term equations, with the Fayet-Iliopoulos terms, become in this case

$$\sum_{b'=1}^{Q_5} (|A_{b'}|^2 - |B_{b'}|^2) = r_3, \quad \sum_{b'=1}^{Q_5} A_{b'} B_{b'} = r_1 + ir_2 \quad (2.89)$$

while the potential is

$$V = \left[\sum_{b'=1}^{Q_5} (|A_{b'}|^2 + |B_{b'}|^2) \right] (\varphi_1^2 + \varphi_2^2 + \varphi_3^2 + \varphi_4^2) \quad (2.90)$$

In this simple case it is easy to see that the presences of the Fayet-Iliopoulos terms in (2.89) ensures that all A, B 's do not vanish simultaneously. The *vev*'s of A, B gives mass to the φ 's. Thus the relative $U(1)$ is broken when the Fayet-Iliopoulos term is not zero. The D-term equations above agree with those in [17] which discusses the splitting of a single D1-brane. It is important to emphasize that the potential and the D-term equations describe an *effective dynamics* in the classical background corresponding to the $(Q_1 - 1, Q_5)$ bound state. This corresponds to the description in [17] of the splitting process in an AdS_3 background which represents a mean field of the above bound state.

⁴The singularity structure for a $U(1)$ theory coupled to N hypermultiplets has been obtained in [23]

2.9.4 The dynamics of the splitting at the singularity of the Higgs branch

We have seen in Section 2.9.2 that the effective theory describing the dynamics of the splitting of the (Q_1, Q_5) system to subystems (Q'_1, Q'_5) and (Q''_1, Q''_5) is $(4, 4)$, $U(1)$ super Yang-Mills coupled to $Q'_1 Q''_5 + Q'_5 Q''_1$ hypermultiplets. The SCFT which this gauge theory flows in the infra-red is singular if the Fayet-Iliopoulos terms and the theta term is set to zero. The description of the superconformal theory of the Higgs branch of a $U(1)$ gauge theory with $(4, 4)$ supersymmetry and N hypermultiplets near the singularity was recently found in [24]. It was expressed in the Coulomb variables. It consists of a bosonic $SU(2)$ Wess-Zumino-Witten model at level $N - 2$, four free fermions and a linear dilaton with background charge given by

$$Q = \sqrt{\frac{2}{N}}(N - 1) \quad (2.91)$$

The central charge of this SCFT is $6(N - 1)$. Using this result for the $U(1)$ theory describing the splitting we get a background charge for the linear dilaton given by

$$Q = \sqrt{\frac{1}{Q'_1 Q''_5 + Q'_5 Q''_1}}(Q'_1 Q''_5 + Q'_5 Q''_1 - 1) \quad (2.92)$$

References

1. J.R. David, G. Mandal and S.R. Wadia, “D1/D5 moduli in SCFT and gauge theory and Hawking radiation,” hep-th/9907075.
2. J. Maldacena, “D-branes and near extremal black holes at low energies,” Phys. Rev. **D55** (1997) 7645.
3. S.F. Hassan and S.R. Wadia, “Gauge theory description of D-brane black holes: emergence of the effective SCFT and Hawking radiation,” Nucl. Phys. **B526** (1998) 311, hep-th/971213.
4. C. Vafa, “Instantons on D-branes,” Nucl. Phys. **B463** (1996) 435, hep-th/9512078.
5. M. R. Douglas, “Branes within branes,” hep-th/9512077.

6. R. Dijkgraaf, “Instanton strings and hyperkaehler geometry,” Nucl. Phys. **B543** (1999) 545, hep-th/9810210.
7. C. Vafa, “Gas of D-branes and Hagedorn Density of BPS States,” Nucl. Phys. **B463** (1996) 415, hep-th/9511088.
8. A. Sen, “U-Duality and Intersecting D-branes,” Phys. Rev. **D53** (1996) 2874, hep-th/9511026.
9. M. Yu, “The unitary representations of the $\mathcal{N} = 4$ $SU(2)$ extended superconformal algebras,” Nucl. Phys. **B294** (1987) 890.
10. L. Dixon, J. Harvey, C. Vafa and E. Witten, “Strings on orbifolds,” Nucl. Phys. **B261** (1985) 620; Strings on orbifolds. 2, Nucl. Phys. **B274** (1986) 285.
11. R. Dijkgraaf, G. Moore, E. Verlinde and H. Verlinde, “Elliptic genera of symmetric products and second quantized strings,” Commun. Math. Phys. **185** (1997) 197, hep-th/9608096.
12. L. Dixon, D. Friedan, E. Martinec and S. Shenker, “The conformal field theory of orbifolds,” Nucl. Phys. **B282** (1987) 13.
13. E. Witten, “Constraints on supersymmetry breaking,” Nucl. Phys. **B202** 253.
14. M. Cvetič, “Effective Lagrangian of the (blown up) orbifolds,” in *Superstrings, Unified Theories And Cosmology* ed. G. Furlan et. al., (1987).
15. S.F. Hassan and S.R. Wadia, “D-brane black holes: large-N limit and the effective string description,” Phys.Lett. **B402** (1997) 43, hep-th/9703163.
16. R. Dijkgraaf, E. Verlinde and H. Verlinde, “5D black holes and matrix strings,” Nucl.Phys. **B506** (1997) 121, hep-th/9704018.
17. N. Seiberg and E. Witten, “The D1/D5 system and singular CFT,” hep-th/9903224.
18. F. Larsen and E. Martinec, “U(1) charges and moduli in the D1/D5 system,” JHEP **9906** (1999) 019, hep-th/9905064.

19. P.S. Aspinwall, “Enhanced gauge symmetries and K_3 surfaces,” Phys. Lett. **B357** (1995) 329, hep-th/9507012.
20. N. Nekrasov and A. Schwarz, “Instantons on noncommutative R^4 and $(2, 0)$ superconformal six dimensional theory,” Commun. Math. Phys. **198** (1998) 689, hep-th/9802068.
21. E. Witten, “Some comments on string dynamics,” hep-th/9507121, in Strings 95, *Future Perspectives In String Theory*, eds. I. Bars et. al.
22. E. Witten, “On the conformal field theory of the Higgs branch,” JHEP **9707**:003 (1997), hep-th/9707093.
23. A. Dhar, G. Mandal, S.R. Wadia and K.P. Yogendran, “D1/D5 system with B-field, noncommutative geometry and the CFT of the Higgs branch,” hep-th/9910194.
24. O. Aharony and M. Berrooz, “IR dynamics of $d = 2, \mathcal{N} = (4, 4)$ gauge theories and DLCQ of ‘Little String Theories’,” hep-th/9909101.

Chapter 3

Coupling with the bulk fields

This chapter focuses on finding the coupling of the microscopic theory discussed in Chapter 2 with fields of the supergravity. The supergravity field ϕ couples with the operator $\mathcal{O}(z, \bar{z})$ of $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ in the form of an interaction

$$S_{\text{int}} = \mu \int d^2z \phi(z, \bar{z}) \mathcal{O}(z, \bar{z}) \quad (3.1)$$

μ denotes the strength of the coupling. We fix the operator $\mathcal{O}(z, \bar{z})$ corresponding to the supergravity field ϕ by appealing to symmetries. We have seen that in Chapter 2 the marginal operators of the SCFT can be classified according to the quantum numbers of the supergroup $SU(1, 1|2) \times SU(1, 1|2)$ together with the $SO(4)_I$ quantum numbers. In this chapter we will see that these symmetries are also present in the near horizon limit of the supergravity solution. We will classify the supergravity fields according to these symmetries. In particular we will focus on all the scalars present in the supergravity background [1]. The coupling in (3.1) is fixed by demanding that the operator $\mathcal{O}(z, \bar{z})$ and the supergravity field ϕ have the same quantum number under the various symmetries [2].

3.1 Near horizon symmetry

In this section we will review the near horizon symmetries of the D1/D5 system. The solution of the type IIB supergravity equations in the string metric is given by [3]

$$\begin{aligned} ds^2 &= f_1^{-\frac{1}{2}} f_5^{-\frac{1}{2}} (-dt^2 + dx_5^2) + f_1^{\frac{1}{2}} f_5^{\frac{1}{2}} (dx_1^2 + \cdots + dx_4^2) \\ &\quad + f_1^{\frac{1}{2}} f_5^{-\frac{1}{2}} (dx_6^2 + \cdots + dx_9^2), \\ e^{-2\phi} &= \frac{1}{g_s^2} f_5 f_1^{-1}, \end{aligned} \quad (3.2)$$

$$\begin{aligned}
B_{05} &= \frac{1}{2}(f_1^{-1} - 1), \\
H_{abc} &= (dB')_{abc} = \frac{1}{2}\epsilon_{abcd}\partial_d f_5, \quad a, b, c, d = 1, 2, 3, 4
\end{aligned}$$

where f_1 and f_5 are given by

$$f_1 = 1 + \frac{16\pi^4 g_s \alpha'^3 Q_1}{V_4 r^2}, \quad f_5 = 1 + \frac{g_s \alpha' Q_5}{r^2}, \quad (3.3)$$

Here $r^2 = x_1^2 + x_2^2 + x_3^2 + x_4^2$ denotes the distance measured in the transverse direction to all the D-branes.

The near horizon scaling limit is obtained by

$$\begin{aligned}
\alpha' &\rightarrow 0, & \frac{r}{\alpha'} &\equiv U = \text{fixed} \\
v &\equiv \frac{V_4}{16\pi^4 \alpha'^2} = \text{fixed}, & g_6 &= \frac{g_s}{\sqrt{v}} = \text{fixed}
\end{aligned} \quad (3.4)$$

In this limit the metric in (3.2) becomes

$$ds^2 = \alpha' \left[\frac{U^2}{g_6 \sqrt{Q_1 Q_5}} (-dx_0^2 + dx_5^2) + g_6 \sqrt{Q_1 Q_5} \frac{dU^2}{U^2} + g_6 \sqrt{Q_1 Q_5} d\Omega_3^2 \right] + \sqrt{\frac{Q_1}{v Q_5}} (dx_6^2 + \dots dx_9^2) \quad (3.5)$$

Thus the near horizon geometry is that of $AdS_3 \times S^3 \times T^4$. Our notation for coordinates here is as follows: $AdS_3 : (x_0, x_5, r); S^3 : (\chi, \theta, \phi); T^4 : (x_6, x_7, x_8, x_9)$. r, χ, θ, ϕ are spherical polar coordinates for the directions x_1, x_2, x_3, x_4 . The radius of S^3 and the anti-de Sitter space is $\sqrt{\alpha'}(g_6^2 Q_1 Q_5)^{1/4}$. Let us examine the symmetries of this near horizon geometry. The bosonic symmetries arise from the isometries of AdS_3 and S^3 . The isometries of the AdS_3 space form the non-compact group $SO(2, 2)$, while the isometries of S^3 form the group $SO(4)_E = SU(2)_E \times \widetilde{SU(2)}_E$. Though the compactification on T^4 breaks the $SO(4)$ rotations of the coordinates $x_6, \dots, x_9$ we can still use this symmetry to classify supergravity fields. We will call this symmetry $SO(4)_I$. The D1/D5 system preserves eight out of the 32 supersymmetries of the type IIB theory. In the near horizon limit the number of supersymmetries gets enhanced from eight to sixteen [4]. These symmetries fix the form of the effective anti-de Sitter supergravity theory near the horizon. The bosonic symmetries $SO(2, 2) \times SO(4)_E = (SL(2R) \times SU(2)) \times (SL(2, R) \times SU(2))$ form the bosonic symmetries of the anti-de Sitter supergravity in three-dimensions. Simple anti-de Sitter supergroups in

three-dimensions were classified in [5]. It can be seen that the only simple supergroups whose bosonic part is $SL(2, R) \times SU(2)$ are $Osp(3|2, R)$ and $SU(1, 1|2)$. The former contains the bosonic subgroup $O(3) \times SL(2, R)$. The supercharges of the supergroup $Osp(3|2, R)$ transform as the vector representation of the group $O(3)$, while the supercharges of the supergroup $SU(1, 1|2)$ transform as **2** of the group $SU(2)$. The unbroken supercharges of the D1/D5 system transform in the spinor representation of $SO(4)_E$ and therefore they transform as **2** of $SU(2)$. This rules out $Osp(3|2, R)$. Therefore the near horizon anti-de Sitter supergravity is based on the supergroup $SU(1, 1|2) \times SU(1, 1|2)$ with matter fields. The pure anti-de Sitter supergravity based on the super group $SU(1, 1|2) \times SU(1, 1|2)$ was constructed in [6] using the fact that it is a Chern-Simons theory .

3.2 Classification of the supergravity modes

In this section we analyze the spectrum of Type IIB supergravity compactified on $AdS_3 \times S^3 \times T^4$. From (3.5) we see the volume of T^4 is $16\pi^4 \alpha'^2 Q_1 / Q_5$. Therefore we ignore Kaluza-Klein modes on the T^4 . The radius of the S^3 is $\sqrt{\alpha'} (g_6 Q_1 Q_5)^{1/4}$. This is large when

$$g_s Q_1 \gg 1 \text{ and } g_s Q_5 \gg 1 \quad (3.6)$$

These inequalities are true in the limit (1.20). Therefore we are justified in using supergravity. Kaluza-Klein reduction of type IIB supergravity to six dimensions leaves six dimensional $(2, 2)$ supergravity. We show that the Kaluza-Klein spectrum of the six dimensional theory on $AdS_3 \times S^3$ can be completely organized as short multiplets of the supergroup $SU(1, 1|2) \times SU(1, 1|2)$. We will follow the method developed by [7].

The massless spectrum of $(2, 2)$ six-dimensional supergravity consists of: a graviton, 8 gravitinos, 5 two-forms, 16 gauge fields, 40 fermions and 25 scalars. Since these are massless, the physical degrees of freedom fall into various representations R_4 of the little group $SO(4)_L$ of $R^{(5,1)}$. For example, the graviton transform as a **(3, 3)** under the little group $SO(4)_L = SU(2)_L \times \widetilde{SU}(2)_L$. On further compactifying $R^{(5,1)}$ into $AdS_3 \times S^3$, each representation R_4 decomposes into various representations R_3 of $SO(3)$, the local Lorentz group of the S^3 . This $SO(3) \simeq SU(2)$ is the diagonal $SU(2)$ of $SU(2)_L \times \widetilde{SU}(2)_L$. For

example, the graviton decomposes as $\mathbf{1} + \mathbf{3} + \mathbf{5}$ under the $SO(3)$, the local Lorentz group of S^3 . The dependence of each of these fields on the angles of S^3 leads to decomposition in terms of Kaluza-Klein modes on the S^3 which transforms according to some representation of the isometry group $SO(4)$ of S^3 . Only those representations of $SO(4)$ occur in these decompositions which contain the representation R_3 of S^3 . To be more explicit, consider the field $\phi_{R_{SO(3)}}(x_0, x_5, r, \theta, \phi, \chi)$ which transforms as some representation $R_{SO(3)}$ of the local Lorentz group of S^3 . The Kaluza-Klein expansion of this field on S^3 is given by

$$\phi_{R_{SO(3)}}(x_0, x_5, r, \theta, \phi, \chi) = \sum_{R_{SO(4)}} \tilde{\phi}_{R_{SO(4)}}(x_0, x_5, r) Y_{R_{SO(3)}}^{R_{SO(4)}}(\theta, \phi, \chi). \quad (3.7)$$

Here $Y_{R_{SO(3)}}^{R_{SO(4)}}(\theta, \phi, \chi)$ stands for the spherical harmonics on S^3 . In the above expansion the only representation of $R_{SO(4)}$ allowed are the ones which contain $R_{SO(3)}$. For example, $\phi(x_0, x_5, r, \theta, \phi, \chi)$ which is a scalar under the local Lorentz group of S^3 can be expanded as

$$\phi(x_0, x_5, r, \theta, \phi, \chi) = \sum_{\mathbf{m}, \mathbf{m}'; \mathbf{m}=\mathbf{m}'} \tilde{\phi}_{\mathbf{m}\mathbf{m}'}(x_0, x_5, r) Y^{(\mathbf{m}, \mathbf{m}')}(\theta, \phi, \chi) \quad (3.8)$$

Once the complete set of Kaluza-Klein modes are obtained we will organize them into short multiplets of the supergroup $SU(1, 1|2) \times SU(1, 1|2)$.

Let us now consider all the massless field of $(2, 2)$ supergravity in six-dimensions individually. The graviton transforms as $(\mathbf{3}, \mathbf{3})$ of the little group in 6 dimensions. The Kaluza-Klein harmonics of this field according to the rules discussed above are

$$\begin{aligned} & (\mathbf{1}, \mathbf{1}) + 2(\mathbf{2}, \mathbf{2}) + (\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}) \\ & + 3 \oplus_{\mathbf{m} \geq \mathbf{3}} (\mathbf{m}, \mathbf{m}) + 2 \oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m} + \mathbf{2}, \mathbf{m}) + (\mathbf{m}, \mathbf{m} + \mathbf{2})] \\ & + \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m} + \mathbf{4}, \mathbf{m}) + (\mathbf{m}, \mathbf{m} + \mathbf{4})] \end{aligned} \quad (3.9)$$

The little group representations of the 8 gravitinos is $4(\mathbf{2}, \mathbf{3}) + 4(\mathbf{3}, \mathbf{2})$. Their Kaluza-Klein harmonics are

$$\begin{aligned} & 8[(\mathbf{1}, \mathbf{2}) + (\mathbf{2}, \mathbf{1})] + 16 \oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m} + \mathbf{1}, \mathbf{m}) + (\mathbf{m}, \mathbf{m} + \mathbf{1})] \\ & + 8 \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m} + \mathbf{3}, \mathbf{m}) + (\mathbf{m}, \mathbf{m} + \mathbf{3})] \end{aligned} \quad (3.10)$$

The Kaluza-Klein harmonics of the 5 two-forms transforming in $(\mathbf{1}, \mathbf{3}) + (\mathbf{3}, \mathbf{1})$ of the little group are

$$10 \oplus_{\mathbf{m} \geq \mathbf{2}} (\mathbf{m}, \mathbf{m}) + 10 \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m} + \mathbf{2}, \mathbf{m}) + (\mathbf{m}, \mathbf{m} + \mathbf{2})] \quad (3.11)$$

The Kaluza-Klein harmonics of the 16 gauge fields, $(\mathbf{2}, \mathbf{2})$ are

$$16(\mathbf{1}, \mathbf{1}) + 32 \oplus_{\mathbf{m} \geq \mathbf{2}} (\mathbf{m}, \mathbf{m}) + 16 \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m}, \mathbf{m} + \mathbf{2}) + (\mathbf{m} + \mathbf{2}, \mathbf{m})] \quad (3.12)$$

The 40 fermions $20(\mathbf{2}, \mathbf{1}) + 20(\mathbf{1}, \mathbf{2})$ give rise to the following harmonics

$$40 \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m}, \mathbf{m} + \mathbf{1}) + (\mathbf{m} + \mathbf{1}, \mathbf{m})] \quad (3.13)$$

The 25 scalars $(\mathbf{1}, \mathbf{1})$ give rise to the harmonics

$$25 \oplus_{\mathbf{m} \geq \mathbf{1}} (\mathbf{m}, \mathbf{m}) \quad (3.14)$$

Putting all this together the complete Kaluza-Klein spectrum of type IIB on $AdS_3 \times S^3 \times T^4$ yields

$$\begin{aligned} &42(\mathbf{1}, \mathbf{1}) + 69(\mathbf{2}, \mathbf{2}) + 48[(\mathbf{1}, \mathbf{2}) + (\mathbf{2}, \mathbf{1})] + 27[(\mathbf{1}, \mathbf{3}) + (\mathbf{3}, \mathbf{1})] \\ &70 \oplus_{\mathbf{m} \geq \mathbf{3}} (\mathbf{m}, \mathbf{m}) + 56 \oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m}, \mathbf{m} + \mathbf{1}) + (\mathbf{m} + \mathbf{1}, \mathbf{m})] \\ &+ 28 \oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m}, \mathbf{m} + \mathbf{2}) + (\mathbf{m} + \mathbf{2}, \mathbf{m})] + 8 \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m}, \mathbf{m} + \mathbf{3}) + (\mathbf{m} + \mathbf{3}, \mathbf{m})] \\ &+ \oplus_{\mathbf{m} \geq \mathbf{1}} [(\mathbf{m}, \mathbf{m} + \mathbf{4}) + (\mathbf{m} + \mathbf{4}, \mathbf{m})] \end{aligned} \quad (3.15)$$

We now organize the above Kaluza-Klein modes into short representations of $SU(1, 1|2) \times SU(1, 1|2)$ [7]. The short multiplet of $SU(1, 1|2)$ consists of the following states

j	L_0	Degeneracy
h	h	$2h + 1$
$h - 1/2$	$h + 1/2$	$2(2h)$
$h - 1$	$h + 1$	$2h - 1$

(3.16)

In the above table j labels the representation of $SU(2)$ which is identified as one of the $SU(2)$'s of the isometry group of S^3 . L_0 denotes the conformal weight of the state. We denote the short multiplet of $SU(1, 1|2) \times SU(1, 1|2)$ as $(\mathbf{2h} + \mathbf{1}, \mathbf{2h}' + \mathbf{1})_S$. On organizing the Kaluza-Klein spectrum into short multiplets we get the following set

$$\begin{aligned} &5(\mathbf{2}, \mathbf{2})_S + 6 \oplus_{\mathbf{m} \geq \mathbf{3}} (\mathbf{m}, \mathbf{m})_S \\ &\oplus_{\mathbf{m} \geq \mathbf{2}} [(\mathbf{m}, \mathbf{m} + \mathbf{2})_S + (\mathbf{m} + \mathbf{2}, \mathbf{m})_S + 4(\mathbf{m}, \mathbf{m} + \mathbf{1})_S + 4(\mathbf{m} + \mathbf{1}, \mathbf{m})_S] \end{aligned} \quad (3.17)$$

Equation (3.15) shows that there are $42(\mathbf{1}, \mathbf{1})$ $SO(4)$ representations in the supergravity Kaluza Klein spectrum. We know that one of these arises from the s -wave of g_{55} from

equation (3.9). This is one of the fixed scalars. $16(\mathbf{1}, \mathbf{1})$ comes from the s -waves of the 16 gauge fields (the components along x^5) as seen in equation (3.12). The remaining 25 comes from the 25 scalars of the six dimensional theory. We would like to see where these $42(\mathbf{1}, \mathbf{1})$ fit in the short multiplets of $SU(1, 1|2) \times SU(1, 1|2)$. From equation (3.17) one can read that 20 of them are in the $5(\mathbf{2}, \mathbf{2})_S$ with $(j = 0, L_0 = 1; j = 0, L_0 = 1)$. 6 of them are in $6(\mathbf{3}, \mathbf{3})_S$ with $(j = 0, L_0 = 2; j = 0, L_0 = 2)$. These correspond to the fixed scalars. Finally, the remaining 16 of them belong to $4(\mathbf{2}, \mathbf{3})_S + 4(\mathbf{3}, \mathbf{2})_S$. 8 of them have $(j = 0, L_0 = 1; j = 0, L_0 = 2)$ and 8 of them have $(j = 0, L_0 = 2; j = 0, L_0 = 1)$. These scalars can be recognized as the intermediate scalars.

3.2.1 Comparison of supergravity shortmultiplets with SCFT

In Section 2.7 we have listed the complete set of shortmultiplets corresponding to single particle states of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$. Comparing equation (2.71) and the list of shortmultiplets of single particle states obtained from supergravity in (3.17) we find that they are identical except for the presence of the following additional shortmultiplets in the SCFT

$$2(\mathbf{1}, \mathbf{2})_S + 2(\mathbf{2}, \mathbf{1})_S + (\mathbf{1}, \mathbf{3})_S + (\mathbf{3}, \mathbf{1})_S \quad (3.18)$$

These correspond to non-propagating degrees of freedom in the supergravity [7]. Therefore they are not present in the list of shortmultiplets obtained from supergravity (3.17).

3.3 The supergravity moduli

In this section we will analyze in detail the massless scalars in the near horizon geometry of the D1/D5 system. Type IIB supergravity compactified on T^4 has 25 scalars. There are 10 scalars h_{ij} which arise from compactification of the metric. $i, j, k \dots$ stands for the directions of T^4 . There are 6 scalars b_{ij} which arise from the Neveu-Schwarz B -field and similarly there are 6 scalars b'_{ij} from the Ramond-Ramond B' -field. The remaining 3 scalars are the ten-dimensional dilaton ϕ_{10} , the Ramond-Ramond scalar χ and the Ramond-Ramond 4-form C_{6789} . These scalars parameterize the coset $SO(5, 5)/(SO(5) \times SO(5))$. The near horizon limit of the D1/D5 system is $AdS_3 \times S^3 \times T^4$ [8]. In this geometry 5 of the 25 scalars become

massive. They are the h_{ii} (the trace of the metric of T^4 which is proportional to the volume of T^4), the 3 components of the anti-self dual part of the Neveu-Schwarz B -field b_{ij}^- and a linear combination of the Ramond-Ramond scalar and the 4-form [9]. The massless scalars in the near horizon geometry parameterize the coset $SO(5, 4)/(SO(5) \times SO(4))$ [10].

As we have seen the near horizon symmetries form the supergroup $SU(1, 1|2) \times SU(1, 1|2)$. We have classified all the massless supergravity fields of type IIB supergravity on $AdS_3 \times S^3 \times T^4$ ignoring the Kaluza-Klein modes on T^4 according to the short multiplets of the supergroup $SU(1, 1|2) \times SU(1, 1|2)$. The isometries of the anti-de Sitter space allow us to relate the quantum number $L_0 + \bar{L}_0$ to the mass of the scalar field through the relation.

$$h + \bar{h} = 1 + \sqrt{1 + m^2} \quad (3.19)$$

Here m is the mass of the scalar in units of the radius of AdS_3 and $(h, \bar{h})$ is the eigenvalue of $L_0, \bar{L}_0$ under the classification of the scalar in shortmultiplets of $SU(1, 1|2) \times SU(1, 1|2)$. Thus the massless fields of the near horizon geometry of the D1/D5 system fall into the top component of the $5(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ short multiplet. We further classify these fields according to the representations of the $SO(4)_I$, the rotations of the x_6, x_7, x_8, x_9 directions. As we have mentioned before this is not a symmetry of the supergravity as it is compactified on T^4 , but it can be used to classify states. The quantum number of the massless supergravity fields are listed below.

Field	$SU(2)_I \times \widetilde{SU}(2)_I$	$SU(2)_E \times \widetilde{SU}(2)_E$	Mass	
$h_{ij} - \frac{1}{4}\delta_{ij}h_{kk}$	$(\mathbf{3}, \mathbf{3})$	$(\mathbf{1}, \mathbf{1})$	0	
b'_{ij}	$(\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \mathbf{3})$	$(\mathbf{1}, \mathbf{1})$	0	
ϕ_6	$(\mathbf{1}, \mathbf{1})$	$(\mathbf{1}, \mathbf{1})$	0	
$a_1\chi + a_2C_{6789}$	$(\mathbf{1}, \mathbf{1})$	$(\mathbf{1}, \mathbf{1})$	0	
b_{ij}^+	$(\mathbf{1}, \mathbf{3})$	$(\mathbf{1}, \mathbf{1})$	0	(3.20)

The linear combination appearing on the fourth line is the one that remains massless in the near-horizon limit. ϕ_6 refers to the six-dimensional dilaton. The $SU(2)_E \times \widetilde{SU}(2)_E$ stands for the $SO(4)$ isometries of the S^3 . All the above fields are s-waves of scalars in the near horizon geometry.

3.4 AdS_3/CFT_2 correspondence

Maldacena's conjecture [12, 13, 14, 15] for the case of the D1/D5 system states that string theory on $AdS_3 \times S^3 \times T^4$ is dual to the 1 + 1 dimensional conformal field theory of the Higgs branch of gauge theory of the D1/D5 system. There is evidence for this conjecture from symmetries. To describe the D1/D5 system at a generic point in the moduli space we can use the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$ to describe the Higgs branch of the gauge theory of the D1/D5 system as we have argued in Chapter 2. The volume of T^4 is of the order of string length and radius of S^3 is large, therefore we can pass over from string theory on $AdS_3 \times S^3 \times T^4$ to six-dimensional (2, 2) supergravity on $AdS_3 \times S^3$. We will compare symmetries in the supergravity limit. The identification of the isometries of the near horizon geometry with that of the symmetries of the SCFT are given in the following table.

Symmetries of the Bulk	Symmetries of SCFT
(a) Isometries of AdS_3 $SO(2, 2) \simeq SL(2, R) \times SL(\widetilde{2}, R)$	The global part of the Virasoro group $SL(2, R) \times SL(\widetilde{2}, R)$
(b) Isometries of S^3 $SO(4)_E \simeq SU(2) \times SU(2)$	R-symmetry of the SCFT $SU(2)_R \times S\widetilde{U}(2)_R$
(c) Sixteen near horizon symmetries	Global supercharges of $\mathcal{N} = (4, 4)$ SCFT
(d) $SO(4)_I$ of T^4	$SO(4)_I$ of $\tilde{T}^4$

To summarize the $SU(1, 1|2) \times SU(1, 1|2)$ symmetry of the near horizon geometry is identified with the global part of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$ together with the identification of the $SO(4)_I$ algebra of T^4 and $\tilde{T}^4$.

3.5 Supergravity moduli and the marginal operators

We would like to match the twenty supergravity moduli appearing in (3.20) with the twenty marginal operators appearing in (2.35) and (2.45) by comparing their symmetry properties under the AdS/CFT correspondence [11].

The symmetries, or equivalently quantum numbers, to be compared under the AdS/CFT

correspondence are as follows:

(a) The isometries of the supergravity are identified with the global symmetries of the superconformal field theory. For the AdS_3 case the symmetries form the supergroup $SU(1,1|2) \times SU(1,1|2)$. The identification of this supergroup with the global part of the $\mathcal{N} = (4,4)$ superalgebra leads to the following mass-dimension relation

$$h + \bar{h} = 1 + \sqrt{1 + m^2} \quad (3.21)$$

where m is the mass of the bulk field and $(h, \bar{h})$ are the dimensions of the SCFT operator. Since in our case the SCFT operators are marginal and the supergravity fields are massless, the mass-dimension relation is obviously satisfied.

(b) The $SU(2)_E \times \widetilde{SU}(2)_E$ quantum number of the bulk supergravity field corresponds to the $SU(2)_R \times \widetilde{SU}(2)_R$ quantum number of the boundary operator. By an inspection of column three of the tables in (2.35), (2.45) and (3.20), we see that these quantum numbers also match.

(c) The location of the bulk fields and the boundary operators as components of the short multiplet can be found by the supersymmetry properties of the bulk fields and the boundary operators. Noting the fact that all the twenty bulk fields as well as all the marginal operators mentioned above correspond to top components of short multiplets, this property also matches.

(d) The above symmetries alone do not distinguish between the twenty operators or the twenty bulk fields. To further distinguish these operators and the fields we identify the $SO(4)_I$ symmetry of the directions x_6, x_7, x_8, x_9 with the $SO(4)_I$ of the SCFT. At the level of classification of states this identification is reasonable though these are not actual symmetries. Using the quantum numbers under this group we obtain the following matching of the boundary operators and the supergravity moduli.

Operator	Field	$SU(2)_I \times \widetilde{SU}(2)_I$	
$\partial x_A^i(z) \bar{\partial} x_A^j(\bar{z}) - 1/4 \delta^{ij} \partial x_A^k \bar{\partial} x_A^k$	$h_{ij} - 1/4 \delta_{ij} h_{kk}$	$(\mathbf{3}, \mathbf{3})$	
$\partial x_A^{[i}(z) \bar{\partial} x_A^{j]}(\bar{z})$	b'_{ij}	$(\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \mathbf{3})$	
$\partial x_A^i(z) \bar{\partial} x_A^i(\bar{z})$	ϕ	$(\mathbf{1}, \mathbf{1})$	
$\mathcal{T}^1$	b_{ij}^+	$(\mathbf{1}, \mathbf{3})$	
$\mathcal{T}^0$	$a_1 \chi + a_2 C_{6789}$	$(\mathbf{1}, \mathbf{1})$	(3.22)

Note that both the representations $(\mathbf{1}, \mathbf{3})$ and $(\mathbf{1}, \mathbf{1})$ occur twice in the above table. This could give rise to a two-fold ambiguity in identifying either $(\mathbf{1}, \mathbf{3})$ or $(\mathbf{1}, \mathbf{1})$ operators with their corresponding bulk fields. The way we have resolved it here is as follows. The operators $\mathcal{T}^1$ and $\mathcal{T}^0$ correspond to blow up modes of the orbifold, and since these are related to the Fayet-Iliopoulos terms and the θ -term in the gauge theory (see Section 2.9), tuning these operators one can reach the singular SCFT [9] that corresponds to fragmentation of the D1/D5 system. In supergravity, similarly, it is only the moduli b_{ij}^+ and $a_1\chi + a_2C_{6789}$ which affect the stability of the D1/D5 system [9, 16, 17]. As a result, it is b_{ij}^+ (and not $b_{ij}'^+$) which should correspond to the operator $\mathcal{T}^1$ and similarly $a_1\chi + a_2C_{6789}$ should correspond to $\mathcal{T}^0$. Another reason for this identification is as follows. b_{ij}^+ and $a_1\chi + a_2C_{6789}$ are odd under world sheet parity while $b_{ij}'^+$ and ϕ are even under world sheet parity. Assign a Z_2 quantum number -1 to the twisted sectors and $+1$ to the untwisted sectors in the SCFT. If under the AdS/CFT correspondence one can identify these Z_2 quantum numbers in the boundary SCFT and the bulk then the correspondence we have made is further justified.

Thus, we arrive at a one-to-one correspondence between operators of the SCFT and the supergravity moduli.

3.6 Fixed scalars

Out of the 25 scalars mentioned earlier which form part of the spectrum of IIB supergravity on T^4 , five become massive when further compactified on $AdS_3 \times S^3$. There is an important additional scalar field which appears after this compactification: h_{55} . Let us remind ourselves the notation used for the coordinates: $AdS_3 : (x_0, x_5, r)$, $S^3 : (\chi, \theta, \phi)$; $T^4 : (x_6, x_7, x_8, x_9)$. r, χ, θ, ϕ are spherical polar coordinates for the directions x_1, x_2, x_3, x_4 . In terms of the D-brane wrappings, the D5 branes are wrapped along the directions x_5, x_6, x_7, x_8, x_9 and D1 branes are aligned along x_5 . The field h_{55} is scalar in the sense that it is a scalar under the local Lorentz group $SO(3)$ of S^3 .

In what follows we will specifically consider the three scalars ϕ_{10} , h_{ii} and h_{55} . The equations of motion of these fields in supergravity are coupled and have been discussed in detail in the literature [18, 19]. It turns out that the six-dimensional dilaton $\phi_6 = \phi_{10} - h_{ii}/4$ which

is a linear combination of h_{ii} and ϕ_{10} remains massless; it is part of the twenty massless (minimal) scalars previously discussed. The two other linear combinations λ and ν are defined as

$$\begin{aligned}\lambda &= \frac{h_{55}}{2} - \frac{\phi_{10}}{2} + \frac{h_{ii}}{8} \\ \nu &= \frac{h_{ii}}{8}\end{aligned}\tag{3.23}$$

λ and ν satisfy coupled differential equations. These are examples of fixed scalars. The pick up masses in the background geometry of the D1/D5 system. In the near horizon limit the equation of motions of λ and ν become decoupled. They give rise to the massive Klein-Gordon equation in AdS_3 . The near horizon mass of λ and ν is $m^2 = 8$ in units of the radius of AdS_3 .

Understanding the absorption and emission properties of fixed scalars is an important problem, because the D-brane computation and semiclassical black hole calculation of these properties appear to be at variance [19]. The discrepancy essentially originates from the ‘expected’ couplings of λ and ν to SCFT operators with $(h, \bar{h}) = (1, 3)$ and $(3, 1)$. These SCFT operators lead to qualitatively different greybody factors from what the fixed scalars exhibit semiclassically. The semiclassical greybody factors are in agreement with D-brane computations if the couplings were only to $(2, 2)$ operators.

The coupling to $(1, 3)$ and $(3, 1)$ operators is guessed from qualitative reasoning based on the Dirac-Born-Infeld action. Since we now have a method of deducing the couplings in (3.1) based on near-horizon symmetries, let us use it to the case of the fixed scalars.

(a) By the mass dimension relation (3.21) we see that the fixed scalars λ and ν correspond to operators with weights $h + \bar{h} = 4$.

(b) The fixed scalars have $SU(2)_E \times \widetilde{SU}(2)_E$ quantum numbers $(\mathbf{1}, \mathbf{1})$.

As all the supergravity fields are classified according to the shortmultiplets of $SU(1, 1|2) \times SU(1, 1|2)$ we can find the field corresponding to these quantum numbers among the shortmultiplets. Searching through the shortmultiplets (see below (3.17)), we find that the fixed scalars belong to the short multiplet $(\mathbf{3}, \mathbf{3})_S$ of $SU(1, 1|2) \times SU(1, 1|2)$. They occur as top component of $(\mathbf{3}, \mathbf{3})_S$. There are six fixed scalars in all. We conclude that the operators with

$(h, \bar{h}) = (1, 3)$ or $(h, \bar{h}) = (3, 1)$ (which were inferred by the DBI method) are ruled out by the analysis of symmetries.

In summary, since the $(1, 3)$ and $(3, 1)$ operators are ruled out by our analysis, the discrepancy between the D-brane calculation and the semiclassical calculation of absorption and emission rates disappears. It is important to note here that couplings guessed from reasoning based on Dirac-Born-Infeld action turns out to be incorrect.

3.7 Intermediate Scalars

We just remark that the classification presented in Section 3.2 correctly account for all sixteen intermediate scalars, and predict that they should couple to SCFT operators with $(h, \bar{h}) = (1, 2)$ belonging to the short multiplet $(\mathbf{2}, \mathbf{3})_S$ or operators with $(h, \bar{h}) = (2, 1)$ belonging to the short multiplet $(\mathbf{3}, \mathbf{2})_S$ (see below equation (3.17)). This agrees with the ‘phenomenological’ prediction made earlier in the literature [20].

3.8 Supergravity from gauge theory

We have seen in this chapter there is a detailed map of the Kaluza-Klein spectrum of super gravity in the near horizon geometry of the D1/D5 system to the shortmultiplets of the $(4, 4)$ SCFT on $\mathcal{M}$. In this section we show that the gauge theory description of the the break up the (Q_1, Q_5) system to subsystems (Q'_1, Q'_5) and Q''_1, Q''_5 agrees with the description of this process in supergravity. In Chapter 2 we have shown that for the (Q_1, Q_5) system to break up into subsystems have to we set the Fayet-Iliopoulos terms and the theta term in the gauge theory to zero. This corresponds to the D1/D5 system with the b_{ij}^+ and $a_1\chi + a_2C_{6789}$ moduli set to zero. From now on we will denote the subsystem (Q'_1, Q'_5) as (Q_1, Q_5) and the subsystem Q''_1, Q''_5 as (q_1, q_5) for convenience. We saw in Section 2.9.4 that in the infrared the description of the splitting process consisted of a linear dilaton with background charge given by

$$Q_{\text{SCFT}} = \sqrt{\frac{2}{Q_1q_5 + Q_5q_1}} (Q_1q_5 + Q_5q_1 - 1) \quad (3.24)$$

We show that the splitting process in supergravity is also controlled by a linear dilaton theory with the background charge given in (3.24).

Lets us first consider the case when $q_5 = 0$ [9]. The Ads/CFT correspondence tells us that we need to consider q_1 D1-branes in the background of $AdS_3 \times S^3 \times T^4$. (We will work in the Euclidean AdS_3 coordinates.) The radius of S^3 and the anti-de Sitter space is given by $r_0 = \sqrt{\alpha'}(g_6^2 Q_1 Q_5)^{1/4}$. For the supergravity to be valid we need to consider the limit (1.20). In the gauge theory the linear dilaton corresponded to the distance between the centre of mass of the two subsystems. Therefore we should focus on the distance between the boundary of $AdS_3 \times S^3 \times T^4$ and the set of q_1 D1-branes. We are interested in the infrared description of the splitting process. By the UV/IR correspondence the D1-branes should be close to the boundary of the $AdS_3 \times S^3 \times T^4$ to obtain the infrared description of the splitting process in the supergravity. We assume that the D1-branes are fixed at a particular point on the S^3 and the T^4 . The action of q_1 D1-branes in the background of AdS_3 and the Ramond-Ramond two-form B_{05} is given by the DBI action. We can use the DBI action for multiple D1-branes as we are interested only in the dynamics of the centre of mass of the collection of q_1 D1-branes. The DBI action of q_1 D1-branes is given by

$$S = \frac{q_1}{2\pi g_s \alpha'} \int d^2 \sigma e^{-\phi} \sqrt{\det (g_{\alpha\beta}^{\text{ind}})} - \frac{q_1}{2\pi g_s \alpha'} \int B \quad (3.25)$$

where σ stands for the world volume coordinates and α, β label these coordinates. $g_{\alpha\beta}^{\text{ind}}$ is the induced metric on the world volume. B is the Ramond-Ramond 2-form potential. We chose a gauge in which the world volume coordinates are the coordinates of the boundary of the AdS_3 . Let the metric on the boundary be $g_{\alpha\beta}(\sigma)$. One can extend the metric $g_{\alpha\beta}(\sigma)$ on the boundary to the interior of AdS_3 in the neighbourhood of the boundary [9]. This is given by

$$ds^2 = \frac{r_0^2}{t^2} \left(dt^2 + \hat{g}_{\alpha\beta}(\sigma, t) d\sigma^\alpha d\sigma^\beta \right) \quad (3.26)$$

with

$$\hat{g}_{\alpha\beta}(\sigma, 0) = g_{\alpha\beta}(\sigma), \quad \hat{g}_{\alpha\beta}(\sigma, t) = g_{\alpha\beta}(\sigma) - t^2 P_{\alpha\beta} + O(t^3) + \dots \quad (3.27)$$

Here $g_{\alpha\beta} P^{\alpha\beta} = R/2$. R is the world sheet curvature. The global coordinates of Euclidean AdS_3 is given by

$$ds^2 = r_0^2 (d\phi^2 + \sinh^2 \phi d\Omega^2) \quad (3.28)$$

where $d\Omega^2$ is the round metric on S^2 . Near the boundary the metric is given by

$$ds^2 = r_0^2(d\phi^2 + \frac{e^{2\phi}}{4}d\Omega^2) \quad (3.29)$$

Motivated by this we use ϕ defined as $t = 2e^{-\phi}$ to measure the distance from the boundary of AdS_3 . Substituting the metric in (3.26) and the near horizon value of the Ramond-Ramond 2-form and the dilaton in (3.25) we obtain the following effective action of the D1-branes near the boundary.

$$\begin{aligned} S &= \frac{q_1 r_0^2}{4\pi g_s \alpha'} \sqrt{\frac{Q_5 v}{Q_1}} \int \sqrt{g} \left(\partial_\alpha \phi \partial^\alpha \phi + \phi R - \frac{1}{2} R + O(e^{-2\phi}) \right) \\ &= \frac{q_1 Q_5}{4\pi} \int \sqrt{g} \left(\partial_\alpha \phi \partial^\alpha \phi + \phi R - \frac{1}{2} R + O(e^{-2\phi}) \right) \end{aligned} \quad (3.30)$$

Now consider the case when $q_1 = 0$. The q_5 D5-branes are wrapped on T^4 . Therefore the world volume of the D5-branes is of the form $M_2 \times T^4$ where M_2 is any 2-manifold. The D5-branes are located at a point on the S^3 . We ignore the fluctuations on T^4 as we are interested in the dynamics on AdS_3 . The DBI action of q_5 D5-branes is given by

$$\frac{q_5}{32\pi^5 g_s \alpha'^3} \left(\int d^6 \sigma e^{-\phi} \sqrt{\det(g_{\alpha\beta}^{\text{ind}})} - \int C^6 \right) \quad (3.31)$$

where C^6 is the Ramond-Ramond 6-form potential coupling to the D5-brane. Performing a similar calculation for the D5-branes and substituting the near horizon values of the 6-form Ramond-Ramond potential, the dilaton and the volume of T^4 one obtains the following effective actions for the D5-branes

$$\begin{aligned} S &= \frac{q_5 r_0^2}{4\pi g_2 \alpha'} \sqrt{\frac{Q_1 v}{Q_5}} \int \sqrt{g} \left(\partial_\alpha \phi \partial^\alpha \phi + \phi R - \frac{1}{2} R + O(e^{-2\phi}) \right) \\ &= \frac{q_5 Q_1}{4\pi} \int \sqrt{g} \left(\partial_\alpha \phi \partial^\alpha \phi + \phi R - \frac{1}{2} R + O(e^{-2\phi}) \right) \end{aligned} \quad (3.32)$$

For the case when $q_1 \neq 0$ and $q_5 \neq 0$ and we just add the contribution from (3.30) and (3.32) to obtain the effective action of the (q_1, q_5) string in AdS_3 . The reason we can do this is because there is no force between the test D1 and D5-branes¹. Thus to the leading order in ϕ the total effective action of the (q_1, q_5) string near the boundary is given by

$$S = \frac{(q_1 Q_5 + q_5 Q_1)}{4\pi} \int \sqrt{g} \left(\partial_\alpha \phi \partial^\alpha \phi + \phi R - \frac{1}{2} R \right) \quad (3.33)$$

¹I thank G. Horowitz and R. Myers for a discussion which helped to clarify this point.

Rescaling ϕ so that the the normalization of the kinetic energy term is canonical one obtains a linear dilaton action with a back ground charge given by

$$Q_{\text{SUGRA}} = \sqrt{2(q_1 Q_5 + q_5 Q_1)} \quad (3.34)$$

We see that the $Q_{\text{SUGRA}} = Q_{\text{SCFT}}$ for large Q_1 and Q_5 . Thus we are able to derive the dynamics of the break up of the (Q_1, Q_5) system to subsystems from the gauge theory.

References

1. J.R. David, G. Mandal and S.R. Wadia, “D1/D5 moduli in SCFT and gauge theory and Hawking radiation,” hep-th/9907075.
2. J.R. David, G. Mandal and S.R. Wadia, “Absorption and Hawking radiation of minimal and fixed scalars, and *AdS/CFT* correspondence,” Nucl. Phys. **B544** (1999) 590, hep-th/9808168.
3. C. G. Callan and J. Maldacena, “D-brane approach to black hole quantum mechanics,” Nucl. Phys. **B472** (1996) 591, hep-th/9602043.
4. P. Claus, R. Kallosh, J. Kumar, P. Townsend and A. Van Proeyen, “Conformal theory of M2, D3, M5 and D1+D5 branes,” JHEP **9806** (1998) 004, hep-th/9801206.
5. M. Günaydin, G. Seirra and P.K. Townsend, “The unitary supermultiplets of d=3 Anti-deSitter and d=2 conformal superalgebras, ” Nucl. Phys. **B274** (1986) 429.
6. J. David, “Anti-de Sitter gravity associated with the supergroup $SU(1, 1|2) \times SU(1, 1|2)$,” Mod. Phys. Lett. **A14** (1999) 1143, hep-th/9904068.
7. J.de Boer, “Six-dimensional supergravity on $S^3 \times AdS_3$ and 2d conformal field theory,” hep-th/9806104.
8. J. Maldacena and A. Strominger, “Black hole greybody factors and D-brane spectroscopy,” Phys. Rev. **D55** (1997) 861, hep-th/9609026; “Universal low-energy dynamics for rotating black holes,” Phys. Rev. **D56** (1997) 4975, hep-th/9702015.

9. N. Seiberg and E. Witten, “The D1/D5 System and singular CFT,” hep-th/9903224.
10. A. Giveon, D. Kutasov and N. Seiberg, “Comments on string theory on AdS_3 ,” Adv. Theor. Math. Phys. **2** (1998) 733, hep-th/9806194.
11. J.R. David, G. Mandal and S.R. Wadia, “Absorption and Hawking radiation of minimal and fixed scalars, and AdS/CFT correspondence,” Nucl. Phys. **B544** (1999) 590, hep-th/9808168.
12. J. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2** (1997) 231, hep-th/9711200.
13. E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2** (1998) 253, hep-th/9802150.
14. S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” hep-th/9802109.
15. J. Maldacena and A. Strominger, “ AdS_3 black holes and a stringy exclusion principle,” JHEP **9812**:005 (1998), hep-th/9804085.
16. F. Larsen and E. Martinec, “U(1) charges and moduli in the D1-D5 system,” JHEP **9906** (1999) 019, hep-th/9905064.
17. R. Dijkgraaf, “Instanton strings and HyperKähler geometry,” Nucl. Phys. **B543** (1999) 545, hep-th/9810210.
18. C. G. Callan, S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “Absorption of fixed scalars and the D-brane approach to black holes,” Nucl. Phys. **B489** (1997) 65, hep-th/9610172.
19. I. R. Klebanov and M. Krasnitz, “Fixed scalar greybody factors in five and four dimensions,” Phys. Rev. **D55** (1997) 3250, hep-th/9612051; “Testing effective string models of black holes with fixed scalars,” Phys. Rev. **D56** (1997) 2173, hep-th/9703216.

20. I. R. Klebanov, A. Rajaraman, A. A. Tseytlin, “Intermediate scalars and the effective string model of black holes,” Nucl. Phys. **B503** (1997) 157, hep-th/9704112.

Chapter 4

The Hawking Process

In this chapter we put the results of Chapter 2 and Chapter 3 together and work towards a precise understanding of Hawking radiation from the D1/D5 black hole starting from the microscopic SCFT [1]. We first review the supergravity calculation of Hawking radiation of minimal scalars from the D1/D5 black hole. We are interested in the D1/D5 black hole with various moduli turned on. We show that in the supergravity calculation the Hawking radiation is independent of the moduli. Then we discuss the SCFT relevant to the D1/D5 black hole and its relationship with the SCFT of the D1/D5 system. We review the formulation of the absorption cross-section calculation from the SCFT as an evaluation of the thermal Green's function of the operators $\mathcal{O}(z, \bar{z})$ corresponding to the supergravity field ϕ . We are able to show that the Hawking radiation calculated from the SCFT agrees precisely with that of the supergravity calculation if we fix the strength of the coupling μ in (1.26) using the AdS/CFT correspondence. We investigate the dependence of the Hawking radiation of minimal scalars calculated from the SCFT and show that it is independent of the moduli as expected from supergravity.

4.1 Supergravity calculation of absorption/Hawking radiation in presence of moduli

We recall that the D1/D5 black hole solution in the absence of moduli is [2, 3] obtained from the D1/D5 system by further compactifying x^5 on a circle of radius R_5 and adding left(right) moving Kaluza-Klein momenta along x^5 . The corresponding supergravity solution is given in (1.14).

The absorption cross-section of minimal scalars in the absence of moduli is given by [4, 5]

$$\sigma_{abs} = 2\pi^2 r_1^2 r_5^2 \frac{\pi\omega}{2} \frac{\exp(\omega/T_H) - 1}{(\exp(\omega/2T_R) - 1)(\exp(\omega/2T_L) - 1)} \quad (4.1)$$

where T_H is given by

$$\frac{2}{T_H} = \frac{1}{T_L} + \frac{1}{T_R} \quad (4.2)$$

The quantities T_L, T_R, r_1, r_5 are as defined in (1.25) and (1.14). We will now show that the absorption cross-section remains unchanged even when the moduli are turned on.

From the equations of motion of type IIB supergravity [6], we can explicitly see that the five-dimensional Einstein metric $ds_{5, Ein}^2$ is not changed by turning on the sixteen moduli corresponding to the metric G_{ij} on T^4 and the Ramond-Ramond 2-form potential B . As regards the four blowing up moduli, the invariance of $ds_{5, Ein}^2$ can be seen from the fact that turning on these moduli corresponds to $SO(4, 5)$ transformation (which is a part of a U-duality transformation) and from the fact that the Einstein metric does not change under U-duality. Now we know that the minimal scalars ϕ^i all satisfy the wave-equation

$$D_\mu \partial^\mu \phi^i = 0 \quad (4.3)$$

where the Laplacian is with respect to the Einstein metric in five dimensions. Since it is only this wave equation that determines the absorption cross-section completely, we see that σ_{abs} is the same as before.

It is straightforward to see that the Hawking rate, given by

$$\Gamma_H = \sigma_{abs} (e^{\omega/T_H} - 1)^{-1} \frac{d^4 k}{(2\pi)^4} \quad (4.4)$$

is also not changed when moduli are turned on.

4.2 Near horizon geometry of the D1/D5 black hole

In order to find out the SCFT relevant for the D1/D5 black hole let us review the near horizon geometry of the D1/D5 black hole. The near horizon scaling limit is given by [7]

$$\alpha' \rightarrow 0, \quad r \rightarrow 0, \quad r_0 \rightarrow 0 \quad (4.5)$$

with

$$\begin{aligned} U &\equiv \frac{r}{\alpha'} = \text{fixed} & U_0 &\equiv \frac{r_0}{\alpha'} = \text{fixed} \\ v &\equiv \frac{V_4}{16\pi^4\alpha'^2} = \text{fixed} & g_6 &= \frac{g_s}{\sqrt{v}} = \text{fixed} \quad R_5 = \text{fixed} \end{aligned} \quad (4.6)$$

In this limit the metric of the D1/D5 black hole (1.14) reduces to the following

$$\begin{aligned} ds^2 &= \frac{\alpha' U^2}{l^2} (-dx_0^2 + dx_5^2) + \frac{\alpha' U_0^2}{l^2} (\cosh \sigma dt + \sinh \sigma dx_5^2)^2 + \frac{\alpha' l^2}{U^2 - U_0^2} dU^2 \\ &+ \alpha' l^2 d\Omega_3^2 + \sqrt{\frac{Q_1}{vQ_5}} (dx_6^2 + \dots dx_9^2) \end{aligned} \quad (4.7)$$

where $l^2 = g_6 \sqrt{Q_1 Q_5}$. It can be seen that this geometry is that of $BTZ \times T^4 \times S^3$ using the co-ordinate redefinitions given below. (Here BTZ refers to the black hole in three-dimensional anti de-Sitter space discovered by [8].)

$$\begin{aligned} \tilde{r}^2 &= (U^2 + U_0^2 \sinh^2 \sigma) \frac{R_5^2}{l^2} \\ r_+ &= \frac{R_5 U_0 \cosh \sigma}{l} \\ r_- &= \frac{R_5 U_0 \sinh \sigma}{l} \\ \phi &= \frac{x_5}{R_5} & t &= \frac{lx_0}{R_5} \end{aligned} \quad (4.8)$$

The metric in these new coordinates is given by

$$\begin{aligned} ds^2 &= -\frac{\alpha' (\tilde{r}^2 - r_+^2)(\tilde{r} - r_-^2)}{l^2 \tilde{r}^2} dt^2 + \frac{\alpha' \tilde{r}^2 l^2}{(\tilde{r} - r_+^2)(\tilde{r} - r_-^2)} dr^2 + \alpha' \tilde{r}^2 \left(d\phi + \frac{r_+ r_-}{\tilde{r}^2 l} dt \right)^2 \\ &+ \alpha' l^2 d\Omega_3^2 + \sqrt{\frac{Q_1}{vQ_5}} (dx_6^2 + \dots + dx_9^2) \end{aligned} \quad (4.9)$$

Our co-ordinate definitions are as follows: $t, \phi, \tilde{r}$ refer to BTZ co-ordinates, Ω_3 stands for the S^3 and x_6, x_7, x_8, x_9 stand for the co-ordinates of T^4 . The mass M and the angular momentum J of the BTZ black hole are given by

$$M = \frac{r_+^2 + r_-^2}{l^2} \quad J = \frac{\sqrt{\alpha'} 2r_+ r_-}{l} \quad (4.10)$$

The mass M and the angular momentum J for the BTZ black hole are related to the parameters of the D1/D5 black hole by

$$\begin{aligned} \frac{M}{2} &= L_0 + \bar{L}_0 = \frac{N_L + N_R}{Q_1 Q_5} \\ \frac{J}{2\sqrt{\alpha'} l} &= L_0 - \bar{L}_0 = \frac{N_L - N_R}{Q_1 Q_5} \end{aligned} \quad (4.11)$$

where N_L , N_R are defined in (1.25) and L_0 , $\bar{L}_0$ are the levels of the SCFT. The extremal limit is given by $r_+ = r_-$. From (4.10) and (4.11) we see that in the extremal limit $N_R = 0$ as expected for the D1/D5 black hole.

In order to find the relevant SCFT corresponding to the D1/D5 black hole let us look at the limit $r_+ = r_- = 0$. Substituting these values in (4.9) we find the metric is given by

$$\frac{ds^2}{\alpha'} = -\frac{\tilde{r}^2}{l^2}dx_0^2 + \frac{l^2}{\tilde{r}^2}dr^2 + \tilde{r}^2d\phi^2 \quad (4.12)$$

In the above equation we have suppressed the T^4 and the S^3 part of the metric. By comparison with (3.5) one can see that this metric is locally AdS_3 except for the global identification $\phi \sim \phi + 2\pi$. Let us compare this metric with that of AdS_3 in this global co-ordinates. In this co-ordinate system the metric of the AdS_3 is given by

$$\frac{ds^2}{\alpha'} = -(\frac{\tilde{r}^2 + l^2}{l^2})dx_0^2 + \frac{l^2}{\tilde{r}^2 + l^2}dt^2 + \tilde{r}^2d\phi^2 \quad (4.13)$$

The geometry of the zero mass BTZ black hole and that of the AdS_3 (although identical locally) has an important difference in the boundary conditions for the fermions. For the case of AdS_3 the fermions are anti-periodic in ϕ and for the zero mass BTZ black hole they are periodic in ϕ . One can easily see that the constant time slice of the metric in (4.13) has the topology of a disk. This forces the fermions to be anti-periodic in ϕ for AdS_3 . For the case of the zero mass BTZ black hole in (4.13), the constant time slice has a singularity at $\tilde{r} = 0$. Therefore the fermions can be both periodic or anti-periodic. An analysis of the killing spinors in the background of the zero mass BTZ [9] shows that the fermions have to be periodic. Thus the SCFT at the boundary of the zero mass BTZ is in the Ramond sector while for the AdS_3 case it is in the Neveu-Schwarz sector.

From the above discussion we find that the SCFT relevant for the D1/D5 black hole with Kaluza-Klein momentum $N = 0$ is the Ramond vacuum of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$. The microscopic states corresponding to the general D1/D5 black hole are states with $L_0 \neq 0$ and $\bar{L}_0 \neq 0$ excited over the Ramond vacuum of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$.

4.3 The coupling with the bulk fields in the Ramond sector

In Chapter 3 the coupling of the supergravity fields ϕ with the operators $\mathcal{O}$ of the $\mathcal{N} = (4, 4)$ SCFT on the $\mathcal{M}$ in (3.1) was determined using the Neveu-Schwarz sector. In this section we argue that these couplings will not change in the Ramond sector. Interaction terms in SCFT Lagrangian do not depend on whether one is in the Ramond sector or in the Neveu-Schwarz sector. It is only the quantities like partition function which change on going from the Neveu-Schwarz sector to the Ramond sector. The scaling dimension of an operator is given by operator product expansions (OPEs) with the stress energy tensor. Since OPEs are local relations, they do not change on going from the Neveu-Schwarz sector to the Ramond sector. Therefore the operators corresponding to the minimal scalars determined in Chapter 3 will not change for the D1/D5 black hole.

4.4 Determination of the strength of the coupling μ

Before we perform the calculation of Hawking radiation/absorption cross-section from the SCFT corresponding to the D1/D5 black hole it is important to determine the strength of the coupling μ in (1.26). In this section we will determine μ for the case of minimal scalars h_{ij} .¹ In Chapter 2 we have identified the SCFT operator corresponding to these fields of the supergravity. The SCFT operator is given by

$$\mathcal{O}^{ij}(z, \bar{z}) = \partial x_A^i(z, \bar{z}) \bar{\partial} x_A^j(z, \bar{z}) - \frac{1}{4} \delta^{ij} \partial x_A^k \bar{\partial} x_A^k(z, \bar{z}) \quad (4.14)$$

Let us suppose the background metric of the torus T^4 is $g_{ij} = \delta_{ij}$. The interaction Lagrangian of the SCFT with the fluctuation h_{ij} is given by

$$S_{\text{int}} = \mu T_{\text{eff}} \int d^2z \left[h_{ij} \partial x_A^i \bar{\partial} x_A^j \right] \quad (4.15)$$

The effective string tension T_{eff} of the conformal field theory, which also appears in the free part of the action

$$S_0 = T_{\text{eff}} \int d^2z \left[\partial_z x_A^i \partial_{\bar{z}} x_{i,A} + \text{fermions} \right] \quad (4.16)$$

¹From now on h_{ij} will denote the traceless part of the metric fluctuations of T^4 .

has been discussed in [6, 12, 13]. The specific value of T_{eff} is not important for the calculation of the S -matrix for absorption or emission, since the factor just determines the normalization of the two-point function of the operator $\mathcal{O}^{ij}(z, \bar{z})$. In this section we will argue that the constant $\mu = 1$.

A direct string theory computation would of course provide the constant μ as well (albeit at weak coupling). This would be analogous to fixing the normalization of the Dirac-Born-Infeld action for a single D-brane by comparing with one-loop open string diagram [16]. However, for a large number and more than one type of D-branes it is a difficult proposition and we will not attempt to pursue it here. Fortunately, the method of symmetries using the AdS/CFT employed for determining the operator $\mathcal{O}$ helps us determine the value of μ as well. For the latter, however, we need to use the more quantitative version [17, 18] of the Maldacena conjecture. We will see below that for this quantitative conjecture to be true for the two-point function (which can be calculated independently from the $\mathcal{N} = (4, 4)$ SCFT and from supergravity) we need $\mu = 1$.

We have seen that the above normalization leads to precise equality between the absorption cross-sections (and consequently Hawking radiation rates) computed from the moduli space of the D1/D5 system and from semiclassical gravity. This method of fixing the normalization can perhaps be criticized on the ground that it borrows from supergravity and does not rely entirely on the SCFT. However, we would like to emphasize two things:

(a) We have fixed $\mu = 1$ by comparing with supergravity around AdS_3 background which does *not* have a black hole. On the other hand, the supergravity calculation of absorption cross-section and Hawking flux is performed around a black hole background represented in the near-horizon limit by the BTZ black hole. From the viewpoint of semiclassical gravity these two backgrounds are rather different. The fact that normalizing μ with respect to the former background leads to the correctly normalized absorption cross-section around the black hole background is a rather remarkable prediction.

(b) Similar issues are involved in fixing the coupling constant between the electron and the electromagnetic field in the semiclassical theory of radiation in terms of the physical electric charge, and in similarly fixing the gravitational coupling of extended objects in terms

of Newton's constant. These issues too are decided by comparing two-point functions of currents with Coulomb's or Newton's laws respectively. In the present case the quantitative version of the AdS/CFT conjecture [17, 18] provides the counterpart of Newton's law or Coulomb's law at strong coupling. Without this the best result one can achieve is that the Hawking radiation rates computed from D1/D5 branes and from semiclassical gravity are *proportional*.

We should remark that fixing the normalization by the use of Dirac-Born-Infeld action, as has been done previously, is not satisfactory since the DBI action is meant for single D-branes and extending it to a system of multiple D1/D5 branes does not always give the right results as we have seen in Section 3.6. The method of equivalence principle to fix the normalization is not very general and cannot be applied to the case of non-minimal scalars, for example.

Let us now compare the two-point function for the minimal scalar h_{ij} determined from the AdS/CFT correspondence and the SCFT to determine the normalization constant μ . We will discuss the more quantitative version of the AdS/CFT conjecture [18, 17] to compare the 2-point correlation function of $\mathcal{O}_{ij}$ from supergravity and SCFT.

The relation between the correlators are as follows. Let the supergravity Lagrangian be

$$\begin{aligned} L &= \int d^3x_1 d^3x_2 b_{ij,i'j'}(x_1, x_2) h_{ij}(x_1) h_{i'j'}(x_2) \\ &+ \int d^3x_1 d^3x_2 d^3x_3 c_{ij,i'j',i''j''}(x_1, x_2, x_3) h_{ij}(x_1) h_{i'j'}(x_2) h_{i''j''}(x_3) + \dots \end{aligned} \quad (4.17)$$

where we have only exhibited terms quadratic and cubic in the h_{ij} 's. The coefficient b determines the propagator and the coefficient c is the tree-level 3-point vertex in supergravity.

The 2-point function of the $\mathcal{O}_{ij}$'s (at large $g_s Q_1, g_s Q_5$) is given by [18, 17], assuming S_{int} given by (4.15)

$$\begin{aligned} &\langle \mathcal{O}_{ij}(z_1) \mathcal{O}_{i'j'}(z_2) \rangle \\ &= 2(\mu T_{\text{eff}})^{-2} \int d^3x_1 d^3x_2 [b_{ij,i'j'}(x_1, x_2) K(x_1|z_1) K(x_2|z_2)] , \end{aligned} \quad (4.18)$$

where K is the boundary-to-bulk Green's function for massless scalars [17].

$$K(x|z) = \frac{1}{\pi} \left[\frac{x_0}{(x_0^2 + (|z_x - z|^2))} \right]^2 \quad (4.19)$$

We use complex z for coordinates of the SCFT, and $x = (x_0, z_x)$ for the Poincaré coordinates of bulk theory.

4.4.1 Evaluation of the tree-level vertices in supergravity

We begin with the bosonic sector of Type IIB supergravity. The Lagrangian is (we follow the conventions of [19])

$$\begin{aligned} I &= I_{\text{NS}} + I_{\text{RR}} \\ I_{\text{NS}} &= -\frac{1}{2k_{10}^2} \int d^{10}x \sqrt{-G} \left[e^{-2\phi} \left(R - 4(d\phi)^2 + \frac{1}{12}(dB_{NS})^2 \right) \right] \\ I_{\text{RR}} &= -\frac{1}{2k_{10}^2} \int d^{10}x \sqrt{-G} \left(\sum_{n=3,7,\dots} \frac{1}{2n!} (H^n)^2 \right) \end{aligned} \quad (4.20)$$

with $k_{10}^2 = 64\pi^7 g_s^2 \alpha'^4$. We use $\hat{M}, \hat{N} \dots$ to denote 10 dimensional indices, $i, j, \dots$ to denote coordinates on the torus T^4 , $M, N \dots$ to denote the remaining 6 dimensions and $\mu, \nu, \dots$ to denote coordinates on the AdS_3 . We have separately indicated the terms depending on Neveu-Schwarz Neveu-Schwarz and Ramond-Ramond backgrounds.

Our aim will be to obtain the Lagrangian of the minimally coupled scalars corresponding to the fluctuations of the metric of the T^4 in the D1/D5-brane system. We will find the Lagrangian up to cubic order in the near horizon limit. Let us first focus on I_{NS} . We substitute the values of the background fields of the D1/D5 system in the Type IIB Lagrangian with the following change in the metric

$$f_1^{\frac{1}{2}} f_5^{-\frac{1}{2}} \delta_{ij} \rightarrow f_1^{\frac{1}{2}} f_5^{-\frac{1}{2}} (\delta_{ij} + h_{ij}). \quad (4.21)$$

where h_{ij} are the minimally coupled scalars with trace zero. These scalars are functions of the 6 dimensional coordinates. Retaining the terms upto $O(h^3)$ and ignoring the traces, the Lagrangian can be written as

$$I_{\text{NS}} = -\frac{V_4}{2k_{10}^2} \int d^6x \sqrt{-G} \frac{G^{MN}}{4} [\partial_M h_{ij} \partial_N h_{ij} + \partial_M (h_{ik} h_{kj}) \partial_N h_{ij}] \quad (4.22)$$

In the above equation we have used the near horizon limit and V_4 is the volume of the T^4 .

The metric G_{MN} near the horizon is

$$ds^2 = \frac{r^2}{R^2}(-dx_0^2 + dx_5^2) + \frac{R^2}{r^2}dr^2 + R^2 d\Omega_3^2 \quad (4.23)$$

We make a change of variables to the Poincaré coordinates by substituting

$$\begin{aligned} z_0 &= \frac{R}{r} \\ z_1 &= \frac{x_0}{R} \\ z_2 &= \frac{x_5}{R} \end{aligned} \quad (4.24)$$

The metric becomes

$$ds^2 = R^2 \frac{1}{z_0^2} (dz_0^2 - dz_1^2 + dz_2^2) + R^2 d\Omega_3^2. \quad (4.25)$$

Here $R = \sqrt{\alpha'}(g_6^2 Q_1 Q_5)^{1/4}$ is the radius of curvature of AdS_3 (also of the S^3). For s -waves the minimal scalars do not depend on the coordinates of the S^3 . Finally, in Poincaré coordinates I_{NS} (correct to cubic order in h) can be written as

$$I_{NS} = -\frac{V_4}{8k_{10}^2} R^3 V_{S^3} \int d^3 z \sqrt{-g} g^{\mu\nu} [\partial_\mu h_{ij} \partial_\nu h_{ij} + \partial_\mu (h_{ik} h_{kj}) \partial_\nu h_{ij}], \quad (4.26)$$

where $V_{S^3} = 2\pi^2$, the volume of a three-sphere of unit radius.

Now we would like to show that to all orders in h , $I_{RR} = 0$ in the near horizon geometry.

The relevant terms in our case are

$$I_{RR} = -\frac{1}{4 \times 3! k_{10}^2} \int d^{10} x \sqrt{-G} H_{\hat{M}\hat{N}\hat{O}} H^{\hat{M}\hat{N}\hat{O}}. \quad (4.27)$$

We substitute the values of B due to the magnetic and electric components of the Ramond-Ramond charges and the value of G . The contribution from the electric part of B' , after going to the near-horizon limit and performing the integral over S^3 and T^4 is

$$\frac{V_4}{4k_{10}^2} R V_{S^3} \int d^3 z \sqrt{-g} \sqrt{\det(\delta_{ij} + h_{ij})} \quad (4.28)$$

The contribution of the magnetic part of B' in the same limit is

$$-\frac{V_4}{4k_{10}^2} R V_{S^3} \int d^3 z \sqrt{-g} \sqrt{\det(\delta_{ij} + h_{ij})} \quad (4.29)$$

We note that the contribution of the electric and the magnetic parts cancel giving no couplings for the minimal scalars to the Ramond-Ramond background. Therefore the tree-level supergravity action correct to cubic order in h is given by

$$I = -\frac{Q_1 Q_5}{16\pi} \int d^3 z [\partial_\mu h_{ij} \partial_\mu h_{ij} + \partial_\mu (h_{ik} h_{kj}) \partial_\mu h_{ij}] \quad (4.30)$$

The coefficient $Q_1 Q_5 / (16\pi)$ is U-duality invariant. This is because it is a function of only the integers Q_1 and Q_5 . This can be tested by computing the same coefficient from the Neveu-Schwarz/fundamental string background which is related to the D1/D5 system by S-duality. The Neveu-Schwarz/fundamental string background also gives the same coefficient. U-duality transformations which generate B_{NS} backgrounds [20] also give rise to the same coefficient.

4.4.2 Two-point function

The two-point function of the operator $\mathcal{O}_{ij}$ can be evaluated by substituting the value of $b_{ij,i'j'}(x_1, x_2)$ obtained from (4.30) into (4.18) and using the boundary-to-bulk Green's function given in (4.19). On evaluating the integral in (4.18) using formulae given in [22], we find that

$$\langle \mathcal{O}_{ij}(z) \mathcal{O}_{i'j'}(w) \rangle = (\mu T_{\text{eff}})^{-2} \delta_{ii'} \delta_{jj'} \frac{Q_1 Q_5}{16\pi^2} \frac{1}{|z - w|^4} \quad (4.31)$$

This is exactly the value of the two-point function obtained from the SCFT described by the free Lagrangian (2.17) provided we put $\mu = 1$.

We have compared the two-point function obtained from the supergravity corresponding to the near horizon geometry of the D1/D5 system with no moduli to the orbifold SCFT. As we have argued before the orbifold SCFT corresponds to the D1/D5 system with moduli. Thus naively this comparison seems to be meaningless. On further examination we note that the coefficient $b_{ij,i'j'}$ in (4.18) was U-duality invariant. Since the D1/D5 system with moduli can be obtained through U-duality transformations we know that this coefficient will not change for the D1/D5 system with moduli. It is only the value of this coefficient which fixes μ to be 1. Thus the comparison we have made is valid. It is remarkable that even at strong coupling the two-point function of $\mathcal{O}_{ij}$ can be computed from the free Lagrangian (4.16). This is consistent with the non-renormalization theorems involving the $\mathcal{N} = (4, 4)$ SCFT.

The choice $\mu = 1$ ensures that the perturbation (4.15) of (4.16) is consistent with the perturbation implied in (4.21). We will see in the next section that this choice leads to precise equality between absorption cross-sections (consequently Hawking radiation rates) calculated from semiclassical gravity and from the D1/D5 branes. The overall multiplicative constant T_{eff} will not be important for the absorption cross-section calculation. This factor finally cancels off in the calculation as we will see in Section 4.7.

4.5 The black hole state

As we have seen, the general non-extremal black hole will have Kaluza-Klein excitations along both the directions on the S^1 . In the SCFT, it is represented by states with $L_0 \neq 0$ and $\bar{L}_0 \neq 0$ over the Ramond vacuum. The black hole is represented by a density matrix

$$\rho = \frac{1}{\Omega} \sum_{\{i\}} |i\rangle\langle i| \quad (4.32)$$

The states $|i\rangle$ belongs to the various twisted sectors of the orbifold theory. They satisfy the constraint

$$L_0 = \frac{N_L}{Q_1 Q_5} \quad \bar{L}_0 = \frac{N_R}{Q_1 Q_5} \quad (4.33)$$

We have suppressed the index which labels the vacuum. Ω is the volume of the phase space in the micro-canonical ensemble. It can be seen that the maximally twisted sector of the orbifold gives rise to the dominant contribution to the sum in (4.32) over the various twisted sectors. The maximally twisted sector is obtained by the action of the twist operator $\sum^{(Q_1 Q_5 - 1)/2}$ on the Ramond vacuum. From the OPE's in (2.55) we see that the twist operator $\sum^{(Q_1 Q_5 - 1)/2}$ introduces a cut in the complex plane such that

$$X_A(e^{2\pi i} z, e^{-2\pi i} \bar{z}) = X_{A+1}(z, \bar{z}) \quad (4.34)$$

Thus this changes the boundary conditions of the bosons and the fermions. Again from the OPEs in (2.55) one infers that the excitations like $\partial\phi_1|\sum^{(Q_1 Q_5 - 1)/2}\rangle$ over the maximally twisted sector have modes in units of $1/Q_1 Q_5$. A simple way of understanding that the maximally twisted sector has modes in units of $1/(Q_1 Q_5)$ is to note that the boundary

conditions in (4.34) imply that $X_A(z, \bar{z})$ is periodic with a period of $2\pi Q_1 Q_5$. This forces the modes to be quantized in units of $1/(Q_1 Q_5)$.

We now show that the maximally twisted sector can account for the entire entropy of the black hole. The entropy of the D1/D5 black hole given in (1.16) can be written as

$$S_{SUGRA} = 2\pi\sqrt{N_L} + 2\pi\sqrt{N_R} \quad (4.35)$$

Using Cardy's formula, the degeneracy of the states in the maximally twisted sector with $L_0 = N_L/Q_1 Q_5$ and $\bar{L}_0 = N_R/Q_1 Q_5$ is given by

$$\Omega = e^{2\pi\sqrt{N_L} + 2\pi\sqrt{N_R}} \quad (4.36)$$

By the Boltzmann formula,

$$S(\text{maximally twisted}) = 2\pi\sqrt{N_L} + 2\pi\sqrt{N_R} \quad (4.37)$$

Thus the maximally twisted sector entirely accounts for the D1/D5 black hole entropy. N_L and N_R are multiples of $Q_1 Q_5$ due to the orbifold projection. Therefore, the entropy can be written as

$$S = 2\pi\sqrt{N_L Q_1 Q_5} + 2\pi\sqrt{N_R Q_1 Q_5} \quad (4.38)$$

With this understanding, we restrict the calculations of Hawking radiation and absorption cross-section only to the maximally twisted sector. The probability amplitude for the Hawking process is given by

$$P = \frac{1}{\Omega} \sum_{f,i} |\langle f | S_{int} | i \rangle|^2 \quad (4.39)$$

where $|f\rangle$ denotes the final states the black hole can decay into. We have averaged over the initial states in the micro-canonical ensemble.

It is more convenient to work with the canonical ensemble. We now discuss the method of determining the temperature of the canonical ensemble. Consider the generating function

$$Z = \text{Tr}_R(e^{-\beta_L E_0} e^{-\beta_R \bar{E}_0}) \quad (4.40)$$

where the trace is done over the Ramond states in the maximally twisted sector. E_0 and $\bar{E}_0$ are energies of the left and the right moving modes.

$$E_0 = \frac{L_0}{R_5}, \quad \bar{E}_0 = \frac{\bar{L}_0}{R_5} \quad (4.41)$$

From the generating function Z in (4.40) we see that the coefficient of $e^{-(\beta_L N_L)/(Q_1 Q_5 R_5)}$ and $e^{-(\beta_R N_R)/(Q_1 Q_5 R_5)}$ is the degeneracy of the states with $L_0 = N_L/Q_1 Q_5$ and $\bar{L}_0 = N_R/Q_1 Q_5$ corresponding to the D1/D5 black hole. A simple way to satisfy this constraint is to choose β_L and β_R such that Z is peaked at this value of L_0 and $\bar{L}_0$.

Evaluating the trace one obtains

$$Z = \prod_{n=1}^{\infty} \left(\frac{1 + e^{-(\beta_L n)/(Q_1 Q_5 R_5)}}{1 - e^{-(\beta_L n)/(Q_1 Q_5 R_5)}} \right)^4 \left(\frac{1 + e^{-(\beta_R n)/(Q_1 Q_5 R_5)}}{1 - e^{-(\beta_R n)/(Q_1 Q_5 R_5)}} \right)^4 \quad (4.42)$$

Then

$$\ln Z = 4 \left[\sum_{n=1}^{\infty} \ln(1 + e^{-\beta_L n/(Q_1 Q_5 R_5)}) - \sum_{n=1}^{\infty} \ln(1 - e^{-\beta_L n/(Q_1 Q_5 R_5)}) \right] \quad (4.43)$$

$$+ 4 \left[\sum_{n=1}^{\infty} \ln(1 + e^{-\beta_R n/(Q_1 Q_5 R_5)}) - \sum_{n=1}^{\infty} \ln(1 - e^{-\beta_R n/(Q_1 Q_5 R_5)}) \right] \quad (4.44)$$

We can evaluate the sum by approximating it by an integral given by

$$\ln Z = 4 Q_1 Q_5 R_5 \int_0^{\infty} dx \ln \left(\frac{1 + e^{-\beta_L x}}{1 - e^{-\beta_L x}} \right) + \ln \left(\frac{1 + e^{-\beta_R x}}{1 - e^{-\beta_R x}} \right) \quad (4.45)$$

From the partition function in (4.40) we see that

$$-\frac{\partial \ln Z}{\partial \beta_L} = \frac{\langle N_L \rangle}{Q_1 Q_5 R_5} \quad \text{and} \quad -\frac{\partial \ln Z}{\partial \beta_R} = \frac{\langle N_R \rangle}{Q_1 Q_5 R_5} \quad (4.46)$$

where $\langle \cdot \rangle$ indicates the average value of N_L and N_R . As the distribution is peaked at N_L and N_R we assume that $\langle N_L \rangle = N_L$ and $\langle N_R \rangle = N_R$. Using (4.45) we obtain

$$\frac{Q_1 Q_5 R_5 \pi^2}{\beta_L^2} = \frac{N_L}{Q_1 Q_5 R_5} \quad \text{and} \quad \frac{Q_1 Q_5 R_5 \pi^2}{\beta_R^2} = \frac{N_R}{Q_1 Q_5 R_5} \quad (4.47)$$

Thus

$$T_L = \frac{1}{\beta_L} = \frac{\sqrt{N_L}}{\pi R_5 Q_1 Q_5} \quad \text{and} \quad T_R = \frac{1}{\beta_R} = \frac{\sqrt{N_R}}{\pi R_5 Q_1 Q_5} \quad (4.48)$$

Above we have introduced a left temperature T_L and a right temperature T_R corresponding to the left and the right moving excitations of the SCFT to pass over to the canonical ensemble.

4.6 Absorption cross-section as thermal Green's function

Let us now relate the absorption cross-section of a supergravity fluctuation $\delta\phi$ to the thermal Green's function of the corresponding operator of the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$ [21]. The notation $\delta\phi$ implies that we are considering the supergravity field to be of the form

$$\phi = \phi_0 + \mu\delta\bar{\phi} \quad (4.49)$$

where ϕ_0 represents the background value and μ is the strength of the coupling.

$$\begin{aligned} S &= S_0 + \int d^2z [\phi_0 + \mu\delta\bar{\phi}] \mathcal{O}(z, \bar{z}) \\ &= S_{\phi_0} + S_{int} \end{aligned} \quad (4.50)$$

where

$$S_{\phi_0} = S_0 + \int d^2z \phi_0 \mathcal{O}(z, \bar{z}) \quad (4.51)$$

$$S_{int} = \mu \int d^2z \delta\bar{\phi} \mathcal{O}(z, \bar{z}) \quad (4.52)$$

$\mathcal{O}$ is the operator corresponding to supergravity field ϕ . S_0 is the Lagrangian of the SCFT which includes the deformations due to various backgrounds in the supergravity. For example, the free Lagrangian in (4.16) corresponds to the case when the field $a_1\chi + a_2C_{6789}$ in (3.20) is turned on.

We calculate the absorption of a quanta $\delta\bar{\phi} = \kappa_5 e^{-ipx}$ corresponding to the operator $\mathcal{O}$ using the Fermi's Golden Rule. κ_5 is related to the five-dimensional Newton's constant G_5 and the ten-dimensional Newton's constant G_{10} as

$$\kappa_5^2 = 8\pi G_5 = \frac{8\pi G_{10}}{V_4 2\pi R_5} = \frac{64\pi^7 g_s^2 \alpha'^4}{V_4 2\pi R_5} \quad (4.53)$$

We see that κ_5 is proportional to α'^2 . In the Maldacena limit (4.5) the coupling of the bulk fluctuation to the SCFT drops out. We retain this term for our calculation as Hawking radiation absorption cross-section is an $O(1/N)$ effect. We will see below that the absorption cross-section turns out to be proportional $g_6 Q_1 Q_5 \alpha'^2$. From (4.7) we see that this is the fourth

power of the radius of S^3 . The radius of S^3 is large for the supergravity to be valid. Thus although in the Maldacena limit (4.5) the coupling of the bulk fluctuation to the SCFT drops out, the factors of $Q_1 Q_5$ picked up in the calculation ensure that this coupling survives for the comparison with supergravity. Furthermore, in analogy with AdS_5/CFT_4 correspondence the number of colours N of the $\mathcal{N} = 4$ SuperYang Mills in four dimensions corresponds to $Q_1 Q_5$. As the absorption cross-section is proportional to $Q_1 Q_5$, this is a $O(1/N)$ effect.

In this computation of the absorption cross-section the black hole is represented by a canonical ensemble at a given temperature. The above interaction gives the thermally averaged transition probability $\mathcal{P}$ as

$$\mathcal{P} = \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} P_{i \rightarrow f} = \mu^2 \kappa_5^2 L t \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} (2\pi)^2 \delta^2(p + p_i - p_f) |\langle f | \mathcal{O}(0,0) | i \rangle|^2 \quad (4.54)$$

Here i and f refer to initial and final states respectively. p_i, p_f refers to the initial and final momenta of these states. $L = 2\pi R_5$ denotes the length of the string and t is the time of interaction. As we have seen in Section 4.5, the inverse temperature β has two components β_L and β_R . The relation of these temperatures to the parameters of the D1/D5 black hole is

$$\beta_L = \frac{1}{T_L} \quad \text{and} \quad \beta_R = \frac{1}{T_R} \quad (4.55)$$

The left moving momenta p_+ and the right moving momenta p_- are in a thermal bath with inverse temperatures β_L and β_R respectively. $\beta \cdot p$ is defined as $\beta \cdot p = \beta_L p_+ + \beta_R p_-$. Z stands for the partition function of the thermal ensemble.

The Green's function in Euclidean time is given by

$$\mathcal{G}(-i\tau, x) = \langle \mathcal{O}^\dagger(-i\tau, x) \mathcal{O}(0,0) \rangle = \text{Tr}(\rho T_\tau \{ \mathcal{O}^\dagger(-i\tau, x) \mathcal{O}(0,0) \}) \quad (4.56)$$

where $\rho = e^{-\beta \cdot \hat{p}}/Z$. Time ordering is defined as T_τ with respect to $-\text{Imaginary}(t)$. This definition coincides with radial ordering on mapping the co-ordinate (τ, x) from the cylinder to the plane. The advantage of doing this is that the integral

$$\int dt dx e^{ip \cdot x} \mathcal{G}(t - i\epsilon, x) = \sum_{i,f} \frac{e^{-\beta \cdot p_i}}{Z} (2\pi)^2 \delta^2(p + p_i - p_f) |\langle f | \mathcal{O}(0,0) | i \rangle|^2 \quad (4.57)$$

The Green's function $\mathcal{G}$ is determined by the two-point function of the operator $\mathcal{O}$. This is in turn determined by conformal dimension $(h, \bar{h})$ of the operator $\mathcal{O}$ and the normalization of the two-point function.

As we have to subtract out the emission probability we get the cross-section as

$$\sigma_{abs} \mathcal{F} t = \mathcal{P}(1 - e^{-\beta \cdot p}) \quad (4.58)$$

where $\mathcal{F}$ is the flux and $\mathcal{P}$ is given by (4.54). Substituting the value of $\mathcal{P}$ from (4.57) we get

$$\sigma_{abs} = \frac{\mu^2 \kappa_5^2 L}{\mathcal{F}} \int dt dx (\mathcal{G}(t - i\epsilon, x) - \mathcal{G}(t + i\epsilon, x)) \quad (4.59)$$

In the above equation we have related the evaluation of the absorption cross-section to the evaluation of the thermal Green's function. Evaluating the integral one obtains

$$\begin{aligned} \sigma_{abs} = & \frac{\mu^2 \kappa_5^2 L \mathcal{C}_{\mathcal{O}} (2\pi T_L)^{2h-1} (2\pi T_R)^{2\bar{h}-1}}{\mathcal{F}} \frac{e^{\beta \cdot p/2} - (-1)^{2h+2\bar{h}} e^{-\beta \cdot p/2}}{2} \\ & \left| \Gamma(h + i \frac{p_+}{2\pi T_L}) \Gamma(\bar{h} + i \frac{p_-}{2\pi T_R}) \right|^2 \end{aligned} \quad (4.60)$$

where $\mathcal{C}_{\mathcal{O}}$ is the coefficient of the leading order term in the OPE of the two-point function of operator $\mathcal{O}$.

4.7 Absorption cross-section of minimal scalars from the D1/D5 SCFT

In the previous section we related the thermal Green's function of the SCFT operator to the absorption cross-section. We will apply the results of the previous section for the case of the minimal scalars. We will consider the case of the minimal scalars corresponding to the fluctuation of the metric of T^4 . Let the background metric of the torus be δ_{ij} . Consider the minimal scalar h_{67} . We know the SCFT operator corresponding to this has conformal dimension $(1, 1)$. From Section 4.4 we know that $\mu = 1$. The interaction Lagrangian is given by

$$S_{int} = 2T_{\text{eff}} \int d^2 z h_{67} \partial x_A^6(z, \bar{z}) \bar{\partial} x_A^7(z, \bar{z}) \quad (4.61)$$

where we have set $\mu = 1$. The factor of 2 arises because of the symmetric property of h_{67} . S_0 is given by

$$S_0 = T_{\text{eff}} \int d^2 z \partial x_A^i(z, \bar{z}) \bar{\partial} x_A^j(z, \bar{z}) \quad (4.62)$$

Comparing with the previous section the operator $\mathcal{O} = 2T_{\text{eff}}\partial x_A^6(z, \bar{z})\bar{\partial}x_A^7(z, \bar{z})$. For the absorption of a quanta of energy ω using (4.60) we obtain

$$\sigma_{abs} = 2\pi^2 r_1^2 r_5^2 \frac{\pi\omega}{2} \frac{\exp(\omega/T_H) - 1}{(\exp(\omega/2T_R) - 1)(\exp(\omega/2T_L) - 1)} \quad (4.63)$$

where we have $L = 2\pi R_5$, $\mathcal{F} = \omega$, (4.53) for κ_5 and (4.31) for $\mathcal{C}_{\mathcal{O}}$. Comparing the absorption cross-section of the minimal scalars obtained from supergravity in (4.1) with (4.63) we find that

$$\sigma_{abs}(\text{ SCFT}) = \sigma_{abs}(\text{ Supergravity}) \quad (4.64)$$

Thus the SCFT calculation and the supergravity calculation of the absorption cross-section agree exactly.

It is important to note that we have used the $\mathcal{N} = (4, 4)$ SCFT realized as a free SCFT on the orbifold $\mathcal{M}$ as the background Lagrangian S_0 . As we have said before, this SCFT is non-singular and therefore cannot correspond to the case of the D1/D5 system with no moduli. In Section 4.1 we have argued that the supergravity calculation of the absorption cross-section is independent of moduli. Therefore it makes sense to compare it with the SCFT result for the case with moduli turned on. In the next section we will show that the SCFT calculation is also independent of the moduli.

Another point worth mentioning is that the method followed in the thesis for the calculation of the absorption-crosssection from the SCFT can be easily extended for the case of minimal scalars corresponding to the four blow up modes. This is not possible if one uses the long string model as the microscopic theory for the D1/D5 black hole. The simple reason being that these operators are not present in the long string model.

4.8 Independence of Hawking radiation on D1/D5 moduli

In this section we will study the independence of the Hawking radiation on D1/D5 moduli. In Chapter 2 we have listed the twenty (1,1) operators $\mathcal{O}_i(z, \bar{z})$ in the SCFT based on the symmetric product orbifold $\mathcal{M}$ which is dual to the D1/D5 system. Turning on various

moduli ϕ^i of supergravity corresponds to perturbing the SCFT

$$S = S_0 + \sum_i \int d^2z \bar{\phi}^i \mathcal{O}_i(z, \bar{z}) \quad (4.65)$$

where $\bar{\phi}^i$ denote the near-horizon limits of the various moduli fields ϕ^i . We note here that S_0 corresponds to the free SCFT based on the symmetric product orbifold $\mathcal{M}$. Since this SCFT is non-singular (all correlation functions are finite), it does not correspond to the marginally stable BPS solution originally found in [2, 3]. Instead, it corresponds to a five-dimensional black hole solution in supergravity with suitable “blow-up” moduli turned on.

Let us now calculate the absorption cross-section of a supergravity fluctuation $\delta\phi_i$ to the thermal Green’s function of the corresponding operator of the SCFT. The notation $\delta\phi_i$ implies that we are considering the supergravity field to be of the form

$$\phi^i = \phi_0^i + \mu\delta\phi^i \quad (4.66)$$

where ϕ_0^i represents the background value and μ is the strength of the coupling.

$$\begin{aligned} S &= S_0 + \int d^2z [\bar{\phi}_0^i + \mu\delta\bar{\phi}^i] \mathcal{O}_i(z, \bar{z}) \\ &= S_{\phi_0} + S_{int} \end{aligned} \quad (4.67)$$

where

$$S_{\phi_0} = S_0 + \int d^2z \bar{\phi}_0^i \mathcal{O}_i(z, \bar{z}) \quad (4.68)$$

$$S_{int} = \mu \int d^2z \delta\bar{\phi}^i \mathcal{O}_i(z, \bar{z}) \quad (4.69)$$

As we have seen in the Section 4.6 the absorption cross-section of the supergravity fluctuation $\delta\phi^i$ involves essentially the two-point function of the operator $\mathcal{O}_i$ calculated with respect to the SCFT action S_{ϕ_0} . Since $\mathcal{O}_i$ is a marginal operator, its two-point function is completely determined apart from a constant. Regarding the marginality of the operators $\mathcal{O}_i$, it is easy to establish it upto one-loop order by direct computation ($c_{ijk} = 0$). The fact that these operators are exactly marginal can be argued as follows. The twenty operators $\mathcal{O}_i$ arise as top components of five chiral primaries. It is known that the number of chiral primaries with $(j_R, \tilde{j}_R) = (m, n)$ is the Hodge number $h_{2m, 2n}$ of the target space $\mathcal{M}$ of the

SCFT. Since this number is a topological invariant, it should be the same at all points of the moduli space of deformations.

We showed in Section 4.4 that if the operator $\mathcal{O}_i$ corresponding to h_{ij} is canonically normalized (OPE has residue 1) and if $\delta\phi_i$ is canonically normalized in supergravity, then the normalization of S_{int} as in (4.69) ensures that σ_{abs} from SCFT agrees with the supergravity result. The crucial point now is the following: once we fix the normalization of S_{int} at a given point in moduli space, at some other point it may acquire a constant ($\neq 1$) in front of the integral when $\mathcal{O}_i$ and $\delta\phi_i$ are canonically normalized at the new point. This would imply that σ_{abs} will get multiplied by this constant, in turn implying disagreement with supergravity. We need to show that this does not happen.

To start with a simple example, let us first restrict to the moduli g_{ij} of the torus $\widetilde{T}^4$. We have

$$S = \int d^2z \partial x^i \bar{\partial} x^j g_{ij} \quad (4.70)$$

The factor of string tension has been absorbed in the definition of x^i .

In Section 4.4 we had $g_{ij} = \delta_{ij} + h_{ij}$, leading to

$$\begin{aligned} S &= S_0 + S_{int} \\ S_0 &= \int d^2z \partial x^i \bar{\partial} x^j \delta_{ij} \\ S_{int} &= \int d^2z \partial x^i \bar{\partial} x^j h_{ij} \end{aligned} \quad (4.71)$$

In the above equation we have set $\mu = 1$. As we have remarked above, this S_{int} gives rise to the correctly normalized σ_{abs} .

Now, if we expand around some other metric

$$g_{ij} = g_{0ij} + h_{ij} \quad (4.72)$$

then the above action (4.70) implies

$$\begin{aligned} S &= S_{g_0} + S_{int} \\ S_{g_0} &= \int d^2z \partial x^i \bar{\partial} x^j g_{0ij} \end{aligned}$$

$$S_{int} = \int d^2z \partial x^i \bar{\partial} x^j h_{ij} \quad (4.73)$$

Now the point is that neither h_{ij} nor the operator $\mathcal{O}^{ij} = \partial X^i \bar{\partial} X^j$ in S_{int} is canonically normalized at $g_{ij} = g_{0ij}$. When we do use the canonically normalized operators, do we pick up an additional constant in front?

Note that

$$\langle \mathcal{O}^{ij} \mathcal{O}^{kl} \rangle_{g_0} = g_0^{ik} g_0^{jl} |z - w|^{-4} \quad (4.74)$$

and

$$\langle h_{ij}(x) h_{kl}(y) \rangle_{g_0} = g_{0,ik} g_{0,jl} \mathcal{D}(x, y) \quad (4.75)$$

where $\mathcal{D}(x, y)$ is the massless scalar propagator

This shows that

Statement (1): *The two-point functions of $\mathcal{O}^{ij}$ and h_{ij} pick up inverse factors.*

As a result, S_{int} remains correctly normalized when re-written in terms of the canonically normalized h and $\mathcal{O}$ and no additional constant is picked up.

The above result is in fact valid in the full twenty dimensional moduli space $\widetilde{\mathcal{M}}$ because Statement (1) above remains true generally.

To see this, let us first rephrase our result for the special case of the metric moduli (4.70) in a more geometric way. The g_{ij} 's can be regarded as some of the coordinates of the moduli space $\widetilde{\mathcal{M}}$ (known to be a coset $SO(4, 5)/(SO(4) \times SO(5))$). The infinitesimal perturbations h_{ij}, h_{kl} can be thought of as defining tangent vectors at the point $g_{0,ij}$ (namely the vectors $\partial/\partial g_{ij}, \partial/\partial g_{kl}$). The (residue of the) two-point function given by (4.74) defines the inner product between these two tangent vectors according to the Zamolodchikov metric [23, 24].

The fact that the moduli space $\widetilde{\mathcal{M}}$ of the $\mathcal{N} = (4, 4)$ SCFT on $\mathcal{M}$ is the coset $SO(4, 5)/(SO(4) \times SO(5))$ is argued in [24]. If the superconformal theory has $\mathcal{N} = (4, 4)$ supersymmetry and if the dimension of the moduli space is d then it is shown in [24] that the moduli space of the SCFT is given by

$$\frac{SO(4, d/4)}{SO(4) \times SO(d/4)} \quad (4.76)$$

The outline of the argument is as follows. An $\mathcal{N} = (4, 4)$ SCFT has superconformal $SU(2)_R \times \widetilde{SU(2)}_R$ symmetry. We have seen that the bottom component of the short multiplet which contains the marginal operator $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ transforms as a $(\mathbf{2}, \mathbf{2})$ under $SU(2)_R \times \widetilde{SU(2)}_R$. The top component which corresponds to the moduli transforms as a $(\mathbf{1}, \mathbf{1})$ under the R-symmetry. The holonomy group of the Zamolodchikov metric should leave invariant the action of $SU(2)_R \times \widetilde{SU(2)}_R$. Then the holonomy group should have a form

$$K \subset SU(2) \times SU(2) \times \tilde{K} \subset SO(d) \quad (4.77)$$

Then (4.77) together with $\mathcal{N} = (4, 4)$ supersymmetry and the left-right symmetry of the two $SU(2)_R$'s of the SCFT fixes the moduli space to be uniquely that given in (4.76). We have found in chapter 2 that there are 20 marginal operators for the $\mathcal{N} = (4, 4)$ SCFT on the orbifold $\mathcal{M}$. Therefore the dimension of the moduli space is 20. Thus $\widetilde{\mathcal{M}}$ is given by

$$\widetilde{\mathcal{M}} = \frac{SO(4, 5)}{SO(4) \times SO(5)} \quad (4.78)$$

Consider, on the other hand, the propagator (inverse two-point function) of h_{ij}, h_{kl} in supergravity. The moduli space action of low energy fluctuations is nothing but the supergravity action evaluated around the classical solutions $g_{0,ij}$. The kinetic term of such a moduli space action defines the metric of moduli space. The statement (1) above is a simple reflection of the fact that the Zamolodchikov metric defines the metric on moduli space, and hence

Statement (2): *The propagator of supergravity fluctuations, viewed as a matrix, is the inverse of the two-point functions in the SCFT.*

The last statement is of course not specific to the moduli g_{ij} and is true of all the moduli. We find, therefore, that fixing the normalization of S_{int} (4.69) at any one point ϕ_0 ensures that the normalization remains correct at any other point ϕ'_0 by virtue of Statement (2). We should note in passing that Statement (2) is consistent with, and could have been derived from AdS/CFT correspondence as applied to the two-point function.

Thus, we find that σ_{abs} is independent of the moduli, in agreement with the result from supergravity.

4.9 Entropy and area

Here we make a brief mention of the fact that the correspondence between Bekenstein-Hawking entropy and the SCFT entropy remains true in the presence of all the twenty moduli. The reason is that in supergravity the Einstein metric remains unchanged (see Section 4.1) and therefore the area of the event horizon remains the same (this can be explicitly verified using the supergravity solution in [20]). In the SCFT, since the operators corresponding to the above moduli are all exactly marginal (Section 4.8) therefore the central charge remains unchanged and hence, by Cardy’s formula, the entropy does not change, in agreement with the Bekenstein-Hawking formula.

References

1. J.R. David, G. Mandal and S.R. Wadia, “Absorption and Hawking radiation of minimal and fixed scalars, and AdS/CFT correspondence,” Nucl. Phys. **B544** (1999) 590, hep-th/9808168.
2. A. Strominger and C. Vafa, “Microscopic origin of the Bekenstein-Hawking entropy,” Phys. Lett. **B379** (1996) 99, hep-th/9601029.
3. C. G. Callan and J. Maldacena, “D-brane approach to black hole quantum mechanics,” Nucl. Phys. **B472** (1996) 591, hep-th/9602043.
4. A. Dhar, G. Mandal and S.R. Wadia, “Absorption vs decay of black holes in string theory and T-symmetry,” Phys. Lett. **B388** (1996) 51, hep-th/9605234.
5. J. Maldacena and A. Strominger, “Black hole greybody factors and D-brane spectroscopy,” Phys. Rev. **D55** (1997) 861, hep-th/9609026.
6. C. G. Callan, S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “Absorption of fixed scalars and the D-brane approach to black holes,” Nucl. Phys. **B489** (1997) 65, hep-th/9610172.

7. J. Maldacena and A. Strominger, “ AdS_3 black holes and a stringy exclusion principle,” JHEP **9812**:005 (1998), hep-th/9804085.
8. M. Banados, C. Teitelboim and J. Zanelli, “Black hole in three-dimensional spacetime,” Phys. Rev. Lett **69** (1992) 1849.
9. O. Coussaert and M. Henneaux, “Supersymmetry of the 2+1 black holes,” Phys. Rev. Lett. **72** (1994) 183, hep-th/9310194.
10. A. Dhar, G. Mandal and S. R. Wadia, “Absorption vs decay of black holes in string theory and T-symmetry,” Phys. Lett. **B388** (1996) 51, hep-th/9605234.
11. S. R. Das, and S. D. Mathur, “Comparing decay rates for black holes and D-branes,” Nucl. Phys. **B478** (1996) 561, hep-th/9606185; “Interactions involving D-branes,” Nucl. Phys. **B482** (1996) 153, hep-th/9607149.
12. Maldacena, “Black holes in string theory,” Ph.D. Thesis, hep-th/9607235.
13. S. F. Hassan and S.R. Wadia, “Gauge theory description of D-brane black holes: emergence of the effective SCFT and Hawking radiation,” Nucl. Phys. **B526** (1998) 311, hep-th/9712213.
14. R. Dijkgraaf, G. Moore, E. Verlinde and H. Verlinde, “Elliptic genera of symmetric products and second quantized strings,” Commun. Math. Phys. **185** (1997) 197, hep-th/9608096.
15. R. Dijkgraaf, E. Verlinde and H. Verlinde, “Matrix string theory,” Nucl. Phys. **B500** (1997) 43, hep-th/9703030.
16. J. Polchinski, S. Chaudhuri and C. V. Johnson, “Notes on D-Branes,” hep-th/9602052; J. Polchinski, “TASI Lectures on D-Branes,” hep-th/9611050.
17. E. Witten, “Anti-de Sitter space and holography,” hep-th/9802150.
18. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” hep-th/9802109.

19. C. P. Bachas, “Lectures on D-branes,” hep-th/9806199.
20. A. Dhar, G. Mandal, S.R. Wadia and K.P. Yogendran, “D1/D5 system with B-field, noncommutative geometry and the CFT of the Higgs branch,” hep-th/9910194.
21. S.S. Gubser, “Absorption of photons and fermions by black holes in four dimensions,” Phys. Rev. **D56** (1997) 7854, hep-th/9706100.
22. Z. Freedman, S. D. Mathur, A. Matusis and L. Rastelli, “Correlation functions in the $\text{CFT}_d/\text{AdS}_{d+1}$ correspondence,” hep-th/9804058.
23. A.A. Zamolodchikov, “Irreversibility of the flux of the renormalization group in a 2-D field theory,” JETP Lett. **43**(1986) 730; Sov. J. Nucl. Phys. 64(1987) 1090.
24. S. Cecotti, “ $\mathcal{N} = 2$ Landau-Ginzburg vs. Calabi-Yau sigma models: nonperturbative aspects,” Int. J. Mod. Phys. **A6** (1991) 1749.

Chapter 5

Concluding remarks and discussions

In this thesis we have made precise the microscopic understanding of Hawking Radiation from the D1/D5 black hole. Our investigations have the following conclusions.

- (a) We presented an explicit construction of all the marginal operators in the SCFT of the D1/D5 system based on the orbifold $\mathcal{M}$. These are twenty in number, four of which are constructed using Z_2 twist operators and correspond to blowing up modes of the orbifold.
- (b) We classified the the twenty near-horizon moduli of supergravity on $\text{AdS}_3 \times S^3 \times T^4$ according to representations of $SU(1,1|2) \times SU(1,1|2) \times SO(4)_I$.
- (c) We established one-to-one correspondence between the supergravity moduli and the marginal operators by inventing a new $SO(4)$ symmetry in the SCFT which we identified with the $SO(4)_I$ of supergravity.
- (d) We have explicitly constructed all the chiral primaries of the $\mathcal{N} = (4,4)$ SCFT on $\mathcal{M}$.
- (e) We analyzed gauge theory dynamics of the D1/D5 system relevant for the splitting of the bound state $(Q_1, Q_5) \rightarrow (Q'_1, Q'_5) + (Q''_1, Q''_5)$.
- (f) We have settled a problem in the context of fixed scalars by showing that consistency with near-horizon symmetry demands that they cannot couple to (1,3) or (3,1) operators. They can only couple to (2,2) operators. This removes earlier discrepancies between D-brane calculations and semiclassical calculations of absorption and emission.
- (g) The black hole is represented by a density matrix of a microcanonical ensemble consisting of states of the various twisted sectors of the $\mathcal{N} = (4,4)$ SCFT on the orbifold $\mathcal{M}$. The dominant contribution to the density matrix is from the maximally twisted sector of the orbifold $\mathcal{M}$. The maximally twisted sector can be obtained by applying the $Q_1 Q_5$ -cycle

twist operator over the Ramond vacuum. The coupling of the bulk field to the orbifold SCFT was deduced using the method of symmetries. These symmetries include the symmetries of the near-horizon geometry, as emphasized by the AdS/CFT correspondence. We calculate the absorption cross-section by relating it to the thermal Greens function of the SCFT. The thermal nature of Hawking radiation is because we have averaged over several microstates.

(h) We determine the normalization of the interaction Lagrangian which couples CFT operators to bulk modes by using the quantitative version of the AdS/CFT correspondence, where we compare two-point functions computed from CFT and from supergravity around AdS_3 background. The normalization fixed this way remarkably leads to precise equality of absorption cross-sections (consequently Hawking radiation rates) computed from CFT and from supergravity around the black hole background.

(i) We showed in supergravity as well as in SCFT that the absorption cross-section for minimal scalars is the same for all values of the moduli, therefore establishing the agreement between SCFT and supergravity all over the moduli space.

We will now discuss some directions for future research.

(a) It is important to determine the absorption cross-section of the fixed scalars from the orbifold SCFT. We have determined all the chiral primaries of the shortmultiplet $(\mathbf{2}, \mathbf{2})_{\mathbf{S}}$ which contains the operators corresponding to the fixed scalars as its top component. We can use these chiral primaries to determine all the operators with conformal dimensions $(h, \bar{h}) = (2, 2)$ corresponding to the fixed scalars. It would be interesting to see if the absorption cross-section calculated using these operators reproduce the semi-classical result.

(b) One can find the description of Hawking radiation of scalars carrying angular momentum within the framework of the SCFT on $\mathcal{M}$.

(d) It is very crucial to find the supergravity solutions of the D1/D5 system with moduli turned on. It is only then we can explicitly check the prediction made in this thesis that Hawking radiation for minimal scalars does not depend on the moduli for the D1/D5 black hole.

(e) Recent developments has suggested that constant B_{NS} moduli is associated with non-commutative geometry. It is important to investigate the D1/D5 system with B_{NS} moduli

turned on. The microscopic theory of the D1/D5 system with B_{NS} moduli encodes information about the moduli space of instantons on a non-commutative torus T^4 .